

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 July 2004 (08.07.2004)

PCT

(10) International Publication Number
WO 2004/056982 A2

(51) International Patent Classification⁷: C12N 9/00

(21) International Application Number:
PCT/EP2003/014674

(22) International Filing Date:
19 December 2003 (19.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0230014.3 23 December 2002 (23.12.2002) GB
60/436,380 23 December 2002 (23.12.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): DEVGEN NV [BE/BE]; Technologiepark 30, B-9052 Gent-Zwijnaarde (BE).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): DE WILDE, Gert, Jules, Hector [BE/BE]; Dr. Armand Rubbensstraat 25, B-9240 Zele (BE).

(74) Agents: MURPHY, Colm, Damien et al.; Boulton Watt, 70 Gray's Inn Road, London WC1X 8BT (GB).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: KINASE SEQUENCES USEFUL FOR DEVELOPING COMPOUNDS FOR THE PREVENTION AND/OR TREATMENT OF METABOLIC DISEASES AND NUCLEOTIDE SEQUENCES ENCODING SUCH KINASE SEQUENCES

(57) Abstract: The present invention relates to nucleotide sequences that encode and may be used to express amino acid sequences that are useful in the identification and development of compounds with activity as pharmaceuticals, in particular of compounds for the prevention and treatment of metabolic diseases such as diabetes and obesity. The invention also relates to the amino acid sequences - such as proteins and polypeptides - that are encoded by, and that may be obtained by suitable expression of, the nucleotide sequences of the invention, particularly the amino acid sequences of J1K, PSK, TAO1 and Q9P2I6. The invention also relates to various uses and modulators of, and methods incorporating, the nucleotide and amino acid sequences of the invention.



WO 2004/056982 A2

Kinase sequences useful for developing compounds for the prevention and/or treatment of metabolic diseases and nucleotide sequences encoding such kinase sequences.

5 The present invention relates to nucleotide sequences that are useful in the pharmaceutical field.

In particular, the invention relates to nucleotide sequences that encode and/or may be used to express amino acid sequences that are useful in the identification and/or development of compounds with (potential) activity as pharmaceuticals, in particular of compounds for the prevention and/or treatment of metabolic diseases such as diabetes
10 and obesity. These nucleotide sequences, which will be further described below, will also be referred to herein as "*nucleotide sequences of the invention*".

The invention also relates to the amino acid sequences - such as proteins and/or polypeptides - that are encoded by, and/or that may be obtained by suitable expression of, the nucleotide sequences of the invention. These amino acid sequences, which will be
15 further described below, will also be referred to herein as "*amino acid sequences of the invention*".

The invention also relates to the use of the nucleotide sequences of the invention, preferably in the form of a suitable genetic construct as described below, in the transformation of host cells and/or host organisms, for example for the expression of the
20 amino acid sequences of the invention. The invention also relates to host cells and/or host organisms that have been transformed with the nucleotide sequences of the invention and/or that can express the amino acid sequences of the invention.

The invention further relates to methods for the identification and/or development of compounds that can modulate the (biological) activity of the amino acid sequences of
25 the invention, in which the abovementioned nucleotide sequences, amino acid sequences, genetic constructs, host cells and/or host organisms are used. Such methods which will usually be in the form of an assay or screen, as will also be further described below.

The invention also relates to the use of the nucleotide sequences, amino acid sequences, genetic constructs, host cells and/or host organisms of the invention in
30 (methods for) identifying and/or developing compounds that can modulate the (biological) activity of the amino acid sequences of the invention.

Also, the invention relates to compounds that can modulate the (biological) activity of the amino acid sequences of the invention, to compositions that contain such compounds, and to the use of such compounds in the preparation of such compositions.

5 In particular, the invention relates to such compositions that are in the form of pharmaceutical compositions, and more in particular in the form of pharmaceutical compositions for the prevention or treatment of metabolic diseases such as diabetes or obesity, and also to the use of compounds that can modulate the (biological) activity of the amino acid sequences of the invention in the preparation of such pharmaceutical compositions.

10 The invention further relates to the use of the nucleotide sequences, amino acid sequences, genetic constructs, host cells and/or host organisms of the invention in (methods for) identifying and/or developing compounds that can be used in the prevention or treatment of metabolic diseases.

Other aspects, embodiments, applications and advantages of the present invention
15 will become clear from the further description below.

The present invention was established from the finding that the amino acid sequences of the invention are involved in metabolic processes (as further described below) and thus can be used as (potential) "target(s)" for *in vitro* and/or *in vivo* interaction with chemical compounds and other factors (with the term "*target*" having its
20 usual meaning in the art, vide for example the definition given in WO 98/06737), and also from the finding that the nucleic acid sequence and amino acid sequences of the invention are involved in metabolic diseases. Consequently, compounds and/or factors that have been identified as interacting with the amino acid sequences of the invention (e.g. by the methods as described hereinbelow) may be useful as active agents in the
25 pharmaceutical field, and in particular for the prevention and treatment of metabolic diseases. All this is supported by the following experimental data/observations:

- In an experimental model for fat handling, metabolism and storage using the model organism *C. elegans* (which model is further described in the Examples), downregulation of the *C. elegans* gene T17E9.1a by RNA interference strongly
30 reduces the fat storage phenotype in said nematode. As further described below, T17E9.1a is the *C. elegans* ortholog of STE20 like kinase in mammals.

Some particularly preferred examples of nucleotide sequences of the invention are:

- the nucleotide sequence of SEQ ID NO:1 (T17E9.1a, also known as *kin-18*), which is a sequence derived from the nematode worm *C. elegans*; and
- the human orthologs of said *C.elegans* sequence, as may be identified by bioinformatic comparison of the *C.elegans* sequence with the human genome. Some preferred, but non-limiting orthologues are given in SEQ ID NOS: 3, 5, 7 and 9.

Generally herein, the use of the human nucleotide sequences of SEQ ID NOS: 3, 5, 7 and 9 and/or the use of nucleotide sequence derived therefrom will be preferred, in particular when the invention is used to develop compounds for pharmaceutical use.

In a broader sense, the term "*nucleotide sequence of the invention*" also comprises:

- parts and/or fragments of any of the nucleotide sequence of SEQ ID NO'S: 1, 3, 5, 7 and 9;
- (natural and/or synthetic) mutants, variants, alleles, analogs, orthologs (hereinbelow collectively referred to as "*mutants*") of any of the nucleotide sequence of SEQ ID NO'S: 1-5, as further described below.
- parts and/or fragments of such (natural or synthetic) mutants;
- nucleotide fusions of any of the nucleotide sequence of SEQ ID NO'S:1-5 (or a part or fragment thereof) with at least one further nucleotide sequence;
- nucleotide fusions of (natural or synthetic) mutants (or a part or fragment thereof) with at least one further nucleotide sequence;

in which such mutants, parts, fragments and/or fusions are preferably as further described below.

The invention also comprises different splice variants of the above nucleotide sequences.

Some particularly preferred examples of amino acid sequences of the invention are:

- the amino acid sequence of SEQ ID NO:2, which is a sequence derived from the nematode worm *C. elegans*; and

- the human analogs of said *C.elegans* sequence, as may be identified by bioinformatic comparison of the *C.elegans* sequence with the human genome. Some preferred, but non-limiting analogs are given in SEQ ID NOS: 4, 6, 8 and 10.

In a broader sense, the term "*amino acid sequence of the invention*" also

5 comprises:

- parts and/or fragments of the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 and 10.
- (natural and/or synthetic) mutants, variants, alleles, analogs, orthologs (hereinbelow collectively referred to as "*analogs*") of the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 and 10;
- 10 - parts and/or fragments of such analogs;
- fusions of the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 and 10 (or a part or fragment thereof) with at least one further amino acid residue or sequence;
- fusions of the amino acid sequence of an analog (or a part or fragment thereof) with at least one further amino acid residue or sequence;

15 in which such mutants, parts, fragments and/or fusions are preferably as further described below.

The term "*amino acid sequence of the invention*" also comprises "immature" forms of the abovementioned amino acid sequences, such as a pre-, pro- or prepro-forms and/or fusions with suitable leader sequences. Also, the amino acid sequences of the
20 invention may have been subjected to post-translational processing and/or be suitably glycosylated, depending upon the host cell or host organism used to express/produce said amino acid sequence; and/or may be otherwise modified (e.g. by chemical techniques known per se in the art).

Generally herein, the use of the human amino acid sequence of SEQ ID NOS: 4,
25 6, 8 and 10 and/or of amino acid sequences derived therefrom will be preferred, in particular when the invention is used to develop compounds for pharmaceutical use.

From the prior art, the following is known about the nucleotide sequences, amino acid sequences and compounds of the invention:

- (1) SEQ ID NOS: 1 and 2. These are the nucleotide sequence and amino acid sequence,
30 respectively, of the *C. elegans* gene/protein T17E9.1a (also known as "*kin-18*")

Reference is made to the following entry from the WORMBASE database:

- for protein sequence:

<http://www.wormbase.org/db/seq/protein?name=WP%3ACE01405;class=Protein>

- for DNA sequence:

<http://www.wormbase.org/db/seq/sequence?name=T17E9.1a;class=Sequence>

5 as well as to Berman et al., Gene 279 (2001) 137-147.

- (2) SEQ ID NOS: 3 and 4. These are the nucleotide sequence and amino acid sequence, respectively, of the human gene/protein known as "*JNK/SAPK inhibitory kinase*" (also known as "*JIK*", "*DPK*" or "*KDS*", or STE20-like kinase).

Reference is made to the following entries from the GENBANK database:

- 10
- NM_016281(mRNA); and
 - NP_057365.2 (translation).

See also:

- GENBANK entry AF135158; Zhang et al., Biochem. Biophys. Res. Commun. 274 (3), 872-879 (2000); MEDLINE 20384190, PUBMED 10924369; and
- 15 - GENBANK entry AF179867; Tassi et al., J. Biol. Chem. 274 (47), 33287-33295 (1999); MEDLINE 20026851; PUBMED 10559204

- (3) SEQ ID NOS: 5 and 6. These are the nucleotide sequence and amino acid sequence, respectively, of the human gene/protein known as "*prostate derived STE20-like kinase*" (also known as "*PSK*" or "*TAO2*").

20 Reference is made to the following entries from the GENBANK database:

- NM_016151.1 (mRNA)
- NP_057235.1 (translation)

See also:

- GENBANK entry AF061943; Moore et al, J. Biol. Chem. 275 (6), 4311-4322
- 25 (2000); MEDLINE 20127920; PUBMED 10660600;
- Chen et al., J. Biol. Chem., Vol. 274, No. 40, pp. 288803-28808 (1999).

- (4) SEQ ID NOS: 7 and 8. These are the nucleotide sequence and amino acid sequence, respectively, of the human gene/protein known as "*thousand and one amino acid protein kinase*" (also known as "*TAO1*").

30 Reference is made to the following entries from the GENBANK database:

- NM_004783 (mRNA)

- NP_004774.1 (translation)

See also:

- GENBANK entry AB020688; Nagase et al, DNA Res. 5 (6), 355-364 (1998);
MEDLINE 99156230; PUBMED 10048485

5 (5) SEQ ID NOS: 9 and 10. These are the nucleotide sequence and amino acid sequence, respectively, of the human gene/protein which with designation KIAA1361 (note that although this gene and protein are also known as "*thousand and one amino acid protein kinase*" or "*TAO1*", KIAA1361 is a different gene/protein from SEQ ID NOS 7 and 8).

10 Reference is made to the following entries from the EMBL and SpTrEMBL databases

- AY049015 (EMBL)
- Q9P2I6 (SPTREMBL)

See also: Nagase et al., DNA Res. 7:65 (2000) and EMBL entry NR AB037782 (<http://www.ebi.ac.uk/cgi-bin/emblfetch?AB037782>)

15 Also, although the inventors do not wish to be limited thereto, it is believed that SEQ ID NOS: 5/6 on the one hand, and SEQ ID NOS: 7 and 8 on the other hand, are transcripts from the same gene/locus.

Thus, in a first aspect, the invention relates to a nucleic acid, preferably in (essentially) isolated form, which nucleic acid encodes and/or can be used to express an amino acid sequence of the invention (as defined herein), and in particular the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 or 10.

In another aspect, the invention relates to a nucleic acid, preferably in (essentially) isolated form, which nucleic acid comprises a nucleotide sequence of the invention, and in particular the nucleotide sequence of SEQ ID NOS: 1, 3, 5, 7 or 9.

25 In a yet another aspect, the invention relates to a nucleic acid, preferably in (essentially) isolated form, which nucleic acid essentially consists of a nucleotide sequence of the invention, and in particular a nucleotide sequence SEQ ID NO: 1, 3, 5, 7 or 9.

Collectively, these nucleic acids will also be referred to herein as "*nucleic acids of the invention*". Also, where appropriate in the context of the further description of the

30

invention below, the terms "*nucleotide sequence of the invention*" and "*nucleic acid of the invention*" may be considered essentially equivalent and essentially interchangeable.

Also, for the purposes of the present invention, a nucleic acid or amino acid sequence is considered to "*(in) essentially isolated (form)*" – for example, from its native
5 biological source - when it has been separated from at least one other component (and in particular macromolecule) with which it is usually associated, such as another nucleic acid, another protein/polypeptide or another (polymeric) biological component. In particular, a nucleic acid or amino acid sequence is considered "essentially isolated" when it has been purified at least 2-fold, in particular at least 10-fold, more in particular
10 at least 100-fold, and up to 1000-fold or more.

The nucleic acids of the invention may also be in the form of a genetic construct, again as further described below. These constructs will also be referred to herein as "*genetic constructs of the invention*". In a preferred embodiment, such a construct will comprise:

- 15 a) the nucleotide sequence of the invention; operably connected to:
b) one or more regulatory elements, such as a promoter and optionally a suitable terminator;
and optionally also:
c) one or more further elements of genetic constructs known per se;
20 in which the terms "*regulatory element*", "*promoter*", "*terminator*", "*further elements*" and "*operably connected*" have the meanings indicated hereinbelow.

In another aspect, the invention relates to a protein or polypeptide, preferably in (essentially) isolated form, said protein or polypeptide comprising an amino acid sequence of the invention (as defined above), and in particular the amino acid sequence
25 of SEQ ID NOS: 2, 4, 6, 8 or 10.

In a further aspect, the invention relates to a protein or polypeptide, preferably in (essentially) isolated form, said protein or polypeptide essentially consisting of an amino acid sequence of the invention (as defined above), and in particular of the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 or 10.

30 In a further aspect, the invention relates to methods for transforming a host cell and/or a host organism with a nucleotide sequence, with a nucleic acid and/or with a

genetic construct of the invention. The invention also relates to the use of a nucleotide sequence, of a nucleic acid and/or of a genetic construct of the invention transforming a host cell or a host organism.

5 In yet another aspect, the invention relates to a host cell or host organism that has been transformed and/or contains with a nucleotide sequence, with a nucleic acid and/or with a genetic construct of the invention. The invention also relates to a host cell and/or host organism that expresses, or (at least) is capable of expressing (e.g. under suitable conditions), an amino acid sequence of the invention. Collectively, such host cells/host organisms will also be referred to herein as "*host cells/host organisms of the invention*".

10 In yet another aspect, the invention relates to methods for producing an amino acid sequence of the invention, in which a nucleotide sequence, nucleic acid, genetic construct, host cell or host organism of the invention is used. Such methods may for instance include expressing a nucleotide sequence of the invention in a suitable host cell or host organism (e.g. upon suitable transformation), and/or maintaining and/or
15 cultivating a host cell or host organism of the invention under suitable conditions, i.e. such that an amino acid sequence of the invention is expressed or obtained. Optionally, these methods may also comprise (one or more steps for) isolating the amino acid sequence thus expressed/produced. The invention also relates to the use of a nucleotide sequence, a nucleic acid, a genetic construct and/or a host cell/host organism of the
20 invention in such a method.

In yet a further aspect, the invention relates to a method for identifying a compound that can modulate the (biological) activity of, and/or that can otherwise interact with, an amino acid sequence of the invention, which method is as further described below. The invention also relates to the use of a nucleotide sequence, a nucleic
25 acid, a genetic construct, an amino acid sequence and/or a host cell/host organism of the invention in such a method.

In yet a further aspect, the invention relates to a method for identifying a compound that can be used in (the preparation of a pharmaceutical composition for) the prevention and/or treatment of metabolic diseases (as further defined below), which
30 method is as further described below. The invention also relates to the use of a nucleotide

sequence, a nucleic acid, a genetic construct, an amino acid sequence and/or a host cell/host organism of the invention in such a method.

The invention also relates to compounds that can modulate the (biological activity of), and/or that can otherwise interact with, an amino acid sequence of the invention,
5 either *in vitro* or preferably (also) *in vivo*, as further described below. The invention also relates to compositions that contain such compounds, and in particular to pharmaceutical compositions that contains such compounds.

The invention further relates to the use of compounds that can modulate the (biological activity of), and/or that can otherwise interact with, an amino acid sequence of
10 the invention in the preparation of these compositions, and in particular to the use of such compounds in the preparation of a pharmaceutical composition for the prevention and/or treatment of metabolic diseases.

The invention also relates to compounds that can be used in the prevention and/or treatment of metabolic diseases (as further defined below), which compounds have or can
15 be identified and/or developed using the method, nucleic acid sequence, amino acid sequence and/or host cell or host organism of the invention. The invention also relates to compositions that contain such compounds, and in particular to pharmaceutical compositions that contain said compounds.

The invention also relates to the use of such compounds in the preparation of a
20 pharmaceutical composition, and in particular to the use of such compounds in the preparation of a pharmaceutical composition for the prevention or treatment of metabolic diseases.

Unless explicitly specified herein, all terms used in the present description have their usual meaning in the art, for which particular reference is made to the definitions
25 given in WO98/06737 and EP 1 085 089.

The nucleotide sequences and amino acid sequences of the invention may generally further be characterized by the presence of a kinase domain (identified using SMART™-analysis), as follows:

30 - SEQ ID NO:2: from amino acid (a.a.) residue 30 to a.a. residue 289

- SEQ ID NO:4: from a.a. residue 24 to a.a. residue 277
- SEQ ID NO:6: from a.a. residue 28 to a.a. residue 281
- SEQ ID NO:8: from a.a. residue 28 to a.a. residue 281
- 5 - SEQ ID NO:10: from a.a. residue 32 to a.a. residue 285

On the basis of the above and on the basis of the prior art referred to above, and although the invention is not specifically limited to any specific explanation or mechanism, it is assumed that the nucleotide sequences and/or amino acid sequences
10 have (biological) activity as kinases.

In particular, and although the invention is again not limited to any specific explanation or hypothesis, it is assumed that the amino acid sequences of the invention may be involved in (the modulation of) the JNK/SAPK pathway (vide for example Zhang et al. and Tassi et al., both supra; as well as Yoneda et al., J. Biol. Chem., Vol. 276,
15 No.17, p. 13935-13940 (2001)).

For JNKs and their role in diabetes/obesity, reference is inter alia made to Hirosumi et al., Nature. Vol. 20, November 2002, 333-336.

As is known in the art, biological activity of this kind can be measured using standard assay techniques for kinases, which are well known to the skilled person. Some
20 preferred, but non-limiting examples include:

- the JNK/SAPK assay described by Zhang et al., supra, which is particularly suitable for the kinase of SEQ ID NO 4 or a similar kinase;
- the kinase assays described by Moore et al., supra, in particular the *in vitro* kinase assays described on page 4315;

25 The nucleotide sequence of SEQ ID NO: 1 was identified, and can be derived/isolated from/using the nematode *C.elegans*; in the manner as further described in Berman et al., or in any other suitable manner known per se.

The nucleotide sequences of SEQ ID NOS: 3, 5, 7 and 9 were identified, and can be derived/isolated from/using human cells; in the manner as further described in the
30 prior art referred to above, or in any other suitable manner known per se.

Also, it is expected that - based upon the disclosure herein - the skilled person will be able to identify, derive and/or isolate natural "mutants" (as mentioned above) of the nucleotide sequence of SEQ ID NOS: 1, 3, 5, 7 and/or 9. For example, such mutants could be derived from (other individuals of) the same species (for example from an individual of a different strain or line, including but not limited to mutant strains or lines); and/or from (individuals of) other species (in which case these mutants will also be referred to herein as "*orthologs*"). Some examples of species from which such orthologs could be derived include, but are not limited to species of

- unicellular and/or micro-organisms such as bacteria, and yeast,
- invertebrate multicellular organisms as such as insects and nematodes (for example, agronomically harmful insect or nematode species);
- vertebrate multicellular organisms as such as fish, birds, reptiles, amphibians and mammals;

Preferably, a natural ortholog is derived from a mammal such as a mouse, rat, rabbit or dog.

Such natural mutants may be obtained in a manner essentially analogous to the methods described in the prior art referred to above, or alternatively by:

- construction of a DNA library from the species of interest in an appropriate expression vector system, followed by direct expression of the mutant sequence;
- construction of a DNA library from the species of interest in an appropriate expression vector system, followed by screening of said library with a probe of the invention (as described below) and/or with a(nother) nucleotide sequence of the invention;
- isolation of mRNA that encodes the mutant sequence from the species of interest, followed by cDNA synthesis using reverse transcriptase;

and/or by any other suitable method(s) or technique(s) known per se, for which reference is for instance made to the standard handbooks, such as Sambrook et al, "Molecular Cloning: A Laboratory Manual" (2nd.ed.), Vols. 1-3, Cold Spring Harbor Laboratory Press (1989) and F. Ausubel et al, eds., "Current protocols in molecular biology", Green Publishing and Wiley Interscience, New York (1987).

It is also expected that - based upon the disclosure herein - the skilled person will be able to provide and/or derive synthetic mutants (as defined hereinabove) of the nucleotide sequences of SEQ ID NOS: 1, 3, 5, 7 and/or 9.

Techniques for generating such synthetic sequences will be clear to the skilled person and may for instance include, but are not limited to, automated DNA synthesis; site-directed mutagenesis; combining two or more parts of one or more naturally occurring sequences, introduction of mutations that lead to the expression of a truncated expression product; introduction of one or more restriction sites (e.g. to create cassettes and/or regions that may easily be digested and/or ligated using suitable restriction enzymes), and/or the introduction of mutations by means of a PCR reaction using one or more "mismatched" primers, using for example a sequence of a naturally occurring GPCR as a template. These and other techniques will be clear to the skilled person, and reference is again made to the standard handbooks, such as Sambrook et al. and Ausubel et al., mentioned above.

Preferably, any mutants as described herein will encode amino acid sequences having one or more, and preferably all, of the structural characteristics/conserved features referred to above for the sequences of SEQ ID NO: 2, 4, 6, 8 and/or 10, and in particular will contain a kinase domain.

It is also possible in the invention to use a part or fragment of the nucleotide sequences of SEQ ID NOS 1, 3, 5, 7 or 9; or a part or fragment of a (natural or synthetic) mutant thereof. These may for instance be 5' and/or 3' truncated nucleotide sequences, or sequences with an introduced *in frame* startcodon or stopcodon. Also, two or more such parts or fragments of one or more nucleotide sequences of the invention may be suitably combined (e.g. ligated in frame) to provide a (further) nucleotide sequence of the invention.

Preferably, any such parts or fragments will be such that they comprise at least one continuous stretch of at least 15 nucleotides, preferably at least 30 nucleotides, more preferably at least 60 nucleotides, even more preferably more than 90 nucleotides, of one or more of the nucleotide sequences of SEQ ID NOS: 1, 3, 5, 7 and/or 9.

In particular, any mutants, parts or fragments as described herein may be such that they (at least) encode the active/catalytic site of the corresponding amino acid sequence

of the invention and/or a binding domain of the corresponding amino acid sequence of the invention

Any mutants, parts and/or fragments as described herein are preferably (also) such that they are capable of hybridizing with one or more of the nucleotide sequences of SEQ ID NOS 1, 3, 5, 7 and/or 9, i.e. under conditions of "moderate stringency", and preferably
5 under conditions of "high stringency". Such conditions will be clear to the skilled person, for example from the standard handbooks, such as Sambrook et al. and Ausubel et al., mentioned above, as well as in EP 0 967 284, EP 1 085 089 or WO 00/55318.

In particular, any mutants, parts and/or fragments as described herein may be such
10 that they are capable of hybridizing with the nucleotide sequence of SEQ ID NO 1, 3, 5, 7 and/or 9 under the "stringent" hybridisation conditions described in WO 00/78972 and WO 98/49185, and/or under the hybridization conditions described in GB 2 357 768-A.

Also, any mutants, parts and/or fragments as described herein will preferably have a degree of "sequence identity", at the nucleotide level, with one or more of the
15 nucleotide sequences of SEQ ID NOS: 1, 3, 5, 7 and/or 9 of at least 50%, preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, and in particular more than 90%, and up to 95% or more.

For this purpose, the percentage of "sequence identity" between a given nucleotide sequence and the nucleotide sequence of SEQ ID NO:1, 3, 5, 7 and/or 9 may
20 be calculated by dividing [*the number of nucleotides in the given nucleotide sequence that are identical to the nucleotide at the corresponding position in the nucleotide sequence of the relevant SEQ ID NO*] by [*the total number of nucleotides in the given nucleotide sequence*] and multiplying by [100%], in which each deletion, insertion, substitution or addition of a nucleotide - compared to the sequence of the relevant SEQ
25 ID NO - is considered as a difference at a single nucleotide (position).

Alternatively, the degree of sequence identity may be calculated using a known computer algorithm for sequence alignment such as NCBI Blast v2.0, using standard settings.

Some other techniques, computer algorithms and settings for determining the
30 degree of sequence identity are for example described in EP 0 967 284, EP 1 085 089, WO 00/55318, WO 00/78972, WO 98/49185 and GB 2 357 768-A.

Also, in a preferred aspect, any mutants, parts and/or fragments as described herein will encode proteins/polypeptides having a biological activity that is essentially similar to the biological activity described above for the sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10, i.e. to a degree of at least 10%, preferably at least 50 % more preferably at least 75%, and up to 90%, as measured by a suitable assay for kinase activity known per se, such as those mentioned above (preferred) or below. In particular, they will have activity as a kinase, as measured using a suitable assay of kinase activity, such as those referred to above, and preferably an activity which is at least 10%, preferably at least 50 % more preferably at least 75%, and up to 90% or more, of the kinase activity of any one of the sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10.

Preferably, any mutants, parts and/or fragments of the nucleotide sequence of the invention will (also) be such that they encode an amino acid sequence which has a degree of "sequence identity", at the amino acid level, with one or more of the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 and/or 10 of at least 50%, preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, and in particular more than 90% and up to 95 % or more, in which the percentage of "sequence identity" is calculated as described below.

Preferably, a nucleotide sequence of the invention will (also) have a length (expressed as total number of nucleotides), which is at least 50%, preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, and in particular more than 90% and up to 95 % or more of the length of one or more of the nucleotide sequence of SEQ ID NOS: 1, 3, 5, 7 and/or 9.

Generally, the nucleotide sequences of the invention, when in the form of a nucleic acid, may be DNA or RNA, and may be single stranded or double stranded. For example, the nucleotide sequences of the invention may be genomic DNA, cDNA or synthetic DNA (such as DNA with a codon usage that has been specifically adapted for expression in the intended host cell or host organism). Thus, the nucleotide sequences of the invention may contain intron sequences, and also generally comprises different splice variants.

It is also within the scope of the invention to use a fusion of a nucleotide sequence of the invention (as described above) with one or more further nucleotide sequence(s),

including but not limited to one or more coding sequences, non-coding sequences and/or regulatory sequences. Preferably, in such fusions, the one or more further nucleotide sequences are operably connected (as described below) to the nucleotide sequence of the invention (for example so that, when the further nucleotide sequence is a coding
5 sequence, the nucleotide fusion encodes a protein fusion as described below).

Another embodiment of the invention relates to a nucleic acid probe that is capable of hybridizing with a nucleotide sequence of the invention under conditions of moderate stringency, preferably under conditions of high stringency, and in particular under stringent conditions (all as described above). Such nucleotide probes may for
10 instance be used for detecting and/or isolating a(nother) nucleotide sequence of the invention and/or as a primer for amplifying a nucleotide sequence of the invention; all using techniques known per se, for which reference is again made to the general handbooks such as Sambrook et al. and Ausubel et al. mentioned above.

Generally, such probes can be designed by the skilled person starting from a
15 nucleotide sequence and/or amino acid sequence of the invention - and in particular one or more of the sequences of SEQ ID NOS 1, 3, 5, 7 and/or 9 - optionally using a suitable computer algorithm. Also, as will be clear to the skilled person, such probes may be degenerate probes.

In another embodiment, the invention relates to an antisense molecule against a
20 nucleotide sequence of the invention.

Yet another embodiment relates to a double stranded RNA molecule directed against a nucleotide sequence of the invention (one strand of which will usually comprise at least part of a nucleotide sequence of the invention). The invention also relates to genetic constructs that can be used to provide such double stranded RNA molecules (e.g.
25 by suitable expression in a host cell or host organism, or for example in a bacterial strain such as *E.coli*). For such constructs, reference is made to for example the International Applications PCT/IB01/1068 and WO 00/01846, both by applicant.

The amino acid sequence of SEQ ID NO: 2 was identified, and can be derived/isolated from/using the nematode *C.elegans*; in the manner as further described
30 in Berman et al., or in any other suitable manner known per se.

The amino acid sequences of SEQ ID NOS: 4, 6, 8 and 10 were identified, and can be derived/isolated from/using human cells; in the manner as further described in the prior art referred to above, or in any other suitable manner known per se.

Generally, the amino acid sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10 may be
5 isolated from the species mentioned above (i.e. *C. elegans* and human, respectively), using any technique(s) for protein isolation and/or purification known per se. Alternatively, the amino acid sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10 may be obtained by suitable expression of a suitable nucleotide sequence - such as one of the nucleotide sequences of SEQ ID NOS: 1, 3, 5, 7 and/or 9, as applicable or a suitable
10 mutant thereof - in an appropriate host cell or host organism, as further described below.

Also, it is expected that - based upon the disclosure herein - the skilled person will be able to identify, derive and/or isolate natural "analogs" (as mentioned above) of the amino acid sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10. Such mutants could be derived from (other individuals of) the same species (for example from an individual of a
15 different strain or line, including but not limited to mutant strains or lines); and/or from (individuals of) other species. For example, such analogs could be derived from the insect species or other pest species mentioned above.

Such natural analogs may again be obtained by isolating them from their natural source using any technique(s) for protein isolation and/or purification known per se, or
20 alternatively by suitable expression of a suitable nucleotide sequence of the invention - such as a natural mutant as described above - in an appropriate host cell or host organism, as further described below.

It is also expected that - based upon the disclosure herein - the skilled person will be able to provide and/or derive synthetic "analogs" (as mentioned above) of one or
25 more of the amino sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10.

Generally, such synthetic analogs may be obtained by suitable expression of a suitable nucleotide sequence of the invention - such as a synthetic mutant as described above - in an appropriate host cell or host organism, as further described below.

Preferably, any analogs as described herein will have one or more, and preferably
30 all, of the structural characteristics/conserved features referred to above for the sequences of SEQ ID NO: 2, 4, 6, 8 and/or 10, and in particular will contain a kinase domain.

It is also possible in the invention to use a part or fragment of one or more of the amino acid sequences of SEQ ID NOS 2, 4, 6, 8 and/or 10, or a part or fragment of a (natural or synthetic) analog thereof mutant thereof. This may for instance be N- and/or C- truncated amino acid sequence. Also, two or more parts or fragments of one or more amino acid sequences of the invention may be suitably combined to provide a (further) amino acid sequence of the invention.

Preferably, any such parts or fragments will be such that they comprise at least one continuous stretch of at least 5 amino acids, preferably at least 10 amino acids, more preferably at least 20 amino acids, even more preferably more than 30 amino acids, of one or more of the amino acid sequences of SEQ ID NO: 2, 4, 6, 8 and/or 10.

In particular, any parts or fragments as described herein are such that they (at least) comprise the active/catalytic site of the corresponding amino acid sequence of the invention and/or a binding domain of the corresponding amino acid sequence of the invention, and in particular a kinase domain. As will be clear to the skilled person, such parts or fragments may find particular use in assay- and screening techniques (as generally described below) and/or (when said part or fragment is provided in crystalline form) in X-ray crystallography.

Generally, such parts or fragments of the amino acid sequences of the invention may be obtained by suitable expression of a suitable nucleotide sequence of the invention - such as a suitable part or fragment as described hereinabove for the nucleotide sequences of the invention - in an appropriate host cell or host organism, as further described below.

In addition and/or as an alternative to the methodology above, amino acid sequences of the invention may also be provided by (chemically and/or enzymatically) modifying the side chain(s) of one or more amino acid residues of an amino acid sequence of SEQ ID NO: 2, 4, 6, 8 and/or 10 or a part, fragment, (natural and/or synthetic) mutant, variant, allele, analogs, orthologs thereof, for example by one or more of the side chain modifications as described in WO 01/02560 and/or by incorporating (e.g. by insertion and/or substitution) one or more unnatural amino acid residues, again as described in WO 01/02560.

Preferably, any analogs, parts and/or fragments as described herein will be such that they have a degree of "sequence identity", at the amino acid level, with one or more of the amino acid sequences of SEQ ID NOS 2, 4, 6, 8 and/or 10 of at least 50%, preferably at least 60%, more preferably at least 70%, even more preferably at least 80%,
5 and in particular more than 90% and up to 95 % or more.

For this purpose, the percentage of "sequence identity" between a given amino acid sequence and one of the amino acid sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10 may be calculated by dividing [*the number of amino acid residues in the given amino acid sequence that are identical to the amino acid residue at the corresponding position*
10 *in the amino acid sequence of the relevant SEQ ID NO*] by [*the total number of amino acid residues in the given amino acid sequence*] and multiplying by [100%], in which each deletion, insertion, substitution or addition of an amino acid residue - compared to the sequence of the relevant SEQ ID NO - is considered as a difference at a single amino acid (position).

15 Alternatively, the degree of sequence identity may be calculated using a known computer algorithm, such as those mentioned above.

Also, such sequence identity at the amino acid level may take into account so-called "conservative amino acid substitutions", which are well known in the art, for example from GB-A-2 357 768, WO 98/49185, WO 00/46383 and WO 01/09300; and
20 (preferred) types and/or combinations of such substitutions may be selected on the basis of the pertinent teachings from the references mentioned in WO 98/49185.

Also, preferably, any analogs, parts and/or fragments as described herein will have a biological activity that is essentially similar to the biological activity described above for the sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10, i.e. to a degree of at least
25 10%, preferably at least 50 %, more preferably at least 75%, and up to 90%, as measured by the assay mentioned above. In particular, they will have activity as a kinase, as measured using a suitable assay of kinase activity, such as those referred to above, and preferably an activity which is at least 10%, preferably at least 50 %, more preferably at least 75%, and up to 90% or more, of the kinase activity of any one of the sequences of
30 SEQ ID NOS: 2, 4, 6, 8 and/or 10.

Preferably, an amino acid sequence of the invention will (also) have a length (expressed as total number of amino acid residues), which is at least 50%, preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, and in particular more than 90% and up to 95 % or more of the length of one or more of the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 and/or 10.

It is also within the scope of the invention to use a fusion of an amino acid sequence of the invention (as described above) with one or more further amino acid sequences, for example to provide a protein fusion. Generally, such fusions may be obtained by suitable expression of a suitable nucleotide sequence of the invention - such as a suitable fusion of a nucleotide sequence of the invention with one or more further coding sequences - in an appropriate host cell or host organism, as further described below. According to one non-limiting embodiment, such a fusion comprises at least the kinase domain of one of the amino acid sequences of the invention.

In one particular embodiment, such fusions may comprise an amino acid sequence of the invention (that preferably at least comprises the kinase domain, and may essentially consist of the kinase domain) fused with a reporter protein such as GFP, luciferase or another fluorescent protein moiety. As will be clear to the skilled person, such fusions may find particular use in expression analysis and similar methodologies.

In another embodiment, the fusion partner may be an amino acid sequence or residue that may be used in purification of the expressed amino acid sequence, for example using affinity techniques directed against said sequence or residue. Thereafter, said sequence or residue may be removed (e.g. by chemical or enzymatical cleavage) to provide the nucleotide sequence of the invention (for this purpose, the sequence or residue may optionally be linked to the amino acid sequence of the invention via a cleavable linker sequence). Some preferred, but non-limiting examples of such residues are multiple histidine residues and glutathione residues,

In one preferred, but non-limiting aspect, any such fusion will have a biological activity that is essentially similar to the biological activity described above for the sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10, i.e. to a degree of at least 10%, preferably at least 50 %, more preferably at least 75%, and up to 90%, as measured by the assay mentioned above. In particular, they will have activity as a kinase, as measured

using a suitable assay of kinase activity, such as those referred to above, and preferably an activity which is at least 10%, preferably at least 50 %, more preferably at least 75%, and up to 90% or more, of the kinase activity of any one of the sequences of SEQ ID NOS: 2, 4, 6, 8 and/or 10.

5 Genetic constructs of the invention will generally comprise at least one nucleotide sequence of the invention, optionally linked to one or more elements of genetic constructs known per se, as described below.

Such genetic constructs may be DNA or RNA, and are preferably double-stranded DNA. The constructs may also be in a form suitable for transformation of the intended
10 host cell or host organism, in a form suitable for integration into the genomic DNA of the intended host cell or in a form suitable for independent replication, maintenance and/or inheritance in the intended host organism. For instance, the genetic construct may be in the form of a vector, such as for example a plasmid, cosmid, YAC, a viral vector or transposon. In particular, the vector may be an expression vector, i.e. a vector that can
15 provide for expression *in vitro* and/or *in vivo* (e.g. in a suitable host cell and/or host organism as described below).

As the one or more "further elements" referred to above, the genetic construct(s) of the invention may generally contain one or more suitable regulatory elements (such as a suitable promoter(s), enhancer(s), terminator(s), etc.), 3'- or 5'-UTR sequences, leader
20 sequences, selection markers, expression markers/reporter genes, and/or elements that may facilitate or increase (the efficiency of) transformation or integration. These and other suitable elements for such genetic constructs will be clear to the skilled person, and may for instance depend upon the type of construct used, the intended host cell or host organism; the manner in which the nucleotide sequences of the invention of interest are to
25 be expressed (e.g. via constitutive, transient or inducible expression); and/or the transformation technique to be used.

Preferably, in the genetic constructs of the invention, the one or more further elements are "*operably linked*" to the nucleotide sequence(s) of the invention and/or to each other, by which is generally meant that they are in a functional relationship with
30 each other. For instance, a promoter is considered "*operably linked*" to a coding sequence if said promoter is able to initiate or otherwise control/regulate the transcription and/or

the expression of a coding sequence (in which said coding sequence should be understood as being "*under the control of*" said promoter)

Generally, when two nucleotide sequences are operably linked, they will be in the same orientation and usually also in the same reading frame. They will usually also be
5 essentially contiguous, although this may also not be required.

Preferably, the optional further elements of the genetic construct(s) used in the invention are such that they are capable of providing their intended biological function in the intended host cell or host organism.

For instance, a promoter, enhancer or terminator should be "*operable*" in the
10 intended host cell or host organism, by which is meant that (for example) said promoter should be capable of initiating or otherwise controlling/regulating the transcription and/or the expression of a nucleotide sequence - e.g. a coding sequence - to which it is operably linked (as defined above).

Such a promoter may be a constitutive promoter or an inducible promoter, and
15 may also be such that it (only) provides for expression in a specific stage of development of the host cell or host organism, and/or such that it (only) provides for expression in a specific cell, tissue, organ or part of a multicellular host organism.

Some particularly preferred promoters include, but are not limited to those present in the expression vectors referred to below.

20 A selection marker should be such that it allows - i.e. under appropriate selection conditions - host cells and/or host organisms that have been (successfully) transformed with the nucleotide sequence of the invention to be distinguished from host cells/organisms that have not been (successfully) transformed. Some preferred, but non-limiting examples of such markers are genes that provide resistance against antibiotics
25 (such as kanamycine or ampicilline), genes that provide for temperature resistance, or genes that allow the host cell or host organism to be maintained in the absence of certain factors, compounds and/or (food) components in the medium that are essential for survival of the non-transformed cells or organisms.

A leader sequence should be such that - in the intended host cell or host organism
30 - it allows for the desired post-translational modifications and/or such that it directs the transcribed mRNA to a desired part or organelle of a cell. A leader sequence may also

allow for secretion of the expression product from said cell. As such, the leader sequence may be any pro-, pre-, or prepro-sequence operable in the host cell or host organism.

An expression marker or reporter gene should be such that - in the host cell or host organism - it allows for detection of the expression of (a gene or nucleotide sequence present on) the genetic construct. An expression marker may optionally also allow for the localisation of the expressed product, e.g. in a specific part or organelle of a cell and/or in (a) specific cell(s), tissue(s), organ(s) or part(s) of a multicellular organism. Such reporter genes may also be expressed as a protein fusion with the amino acid sequence of the invention. Some preferred, but non-limiting examples include fluorescent proteins such as Green Fluorescent Protein (GFP).

For some (further) non-limiting examples of the promoters, selection markers, leader sequences, expression markers and further elements that may be present/used in the genetic constructs of the invention - such as terminators, transcriptional and/or translational enhancers and/or integration factors - reference is made to the general handbooks such as Sambrook et al. and Ausubel et al. mentioned above, to W.B. Wood et al., "*The nematode Caenorhabditis elegans*", Cold Spring Harbor Laboratory Press (1988) and D.L. Riddle et al., "*C. ELEGANS II*", Cold Spring Harbor Laboratory Press (1997), as well as to the examples that are given in WO 95/07463, WO 96/23810, WO 95/07463, WO 95/21191, WO 97/11094, WO 97/42320, WO 98/06737, WO 98/21355, US-A-6,207,410, US-A- 5,693,492 and EP 1 085 089. Other examples will be clear to the skilled person.

The genetic constructs of the invention may generally be provided by suitably linking the nucleotide sequence(s) of the invention to the one or more further elements described above, for example using the techniques described in the general handbooks such as Sambrook et al. and Ausubel et al., mentioned above.

Often, the genetic constructs of the invention will be obtained by inserting a nucleotide sequence of the invention in a suitable (expression) vector known per se. Some preferred, but non-limiting examples of suitable expression vectors include:

- vectors for expression in mammalian cells: pMAMneo (Clontech), pcDNA3 (Invitrogen), pMC1neo (Stratagene), pSG5 (Stratagene), EBO-pSV2-neo (ATCC 37593), pBPV-1 (8-2) (ATCC 37110), pdBPV-MMTneo (342-12) (ATCC 37224),

pRSVgpt (ATCC37199), pRSVneo (ATCC37198), pSV2-dhfr (ATCC 37146),
pUCTag (ATCC 37460) and 1ZD35 (ATCC 37565);

- vectors for expression in bacterial cells: pET vectors (Novagen) and pQE vectors (Qiagen);
- 5 - vectors for expression in yeast or other fungal cells: pYES2 (Invitrogen) and Pichia expression vectors (Invitrogen);
- vectors for expression in insect cells: pBlueBacII (Invitrogen).

The nucleotide sequences and/or genetic constructs of the invention may be used to transform a host cell or host organism.

10 The host cell may be any suitable (fungal, prokaryotic or eukaryotic) cell or cell line, for example:

- a bacterial strain, including but not limited to strains of *E.coli*, *Bacillus*, *Streptomyces* and *Pseudomonas*;
- a fungal cell, including but not limited to cells from species of *Aspergillus* and
15 *Trichoderma*;
- a yeast cell, including but not limited to cells from species of *Kluyveromyces* or *Saccharomyces*;
- an amphibian cell or cell line, such as *Xenopus* oocytes.

In one specific embodiment, which may prove particularly useful when the
20 nucleotide sequences of the invention are (to be) used in the discovery and development of insecticidal compounds, the host cell may be an insect-derived cell or cell line, such as:

- cells/cell lines derived from *lepidoptera*, including but not limited to *Spodoptera* SF9 and Sf21 cells,
- 25 - cells/cell lines derived from *Drosophila*, such as Schneider and Kc cells; and/or
- cells/cell lines derived from a pest species of interest (as mentioned below), such as from *Heliothis virescens*.

In one preferred embodiment, the host cell is a mammalian cell or cell line, for example derived from the mammals referred to above.

In an even more preferred aspect, the host cell is a cell or cell line derived from a human, and from other mammals including but not limited to CHO- and BHK-cells and human cells or cell lines such as HeLa and COS.

In one specific, but non-limiting embodiment, the cell or cell line may be a human
5 cell or cell line which is related to metabolic processes or metabolic disease and/or used as a cellular model for metabolic disease, including but not limited to liver cells or cell lines, adipocytes or muscle cells or cell lines such as HEPG2 cells, 3T3L1 adipocytes, CTC12 cells and L6 myotubes.

The host organism may be any suitable multicellular (vertebrate or invertebrate)
10 organism, including but not limited to:

- a nematode, including but not limited to nematodes from the genus *Caenorhabditis*, such as *C.elegans*,
- an insect, including but not limited to species of *Drosophila* and/or a specific pest species of interest (such as those mentioned above);
- 15 - other well known model organisms, such as zebrafish;
- a mammal such as a rat or mouse;

Other suitable host cells or host organisms will be clear to the skilled person, for example from the handbooks and patent applications mentioned above.

It should be noted that when a nucleotide sequence of the invention is expressed
20 in a multicellular organism, it may be expressed throughout the entire organism, or only in one or more specific cells, tissues, organs and/or parts thereof, for example by expression under the control of a promoter that is specific for said cell(s), tissue(s), organ(s) or part(s).

The nucleotide sequence may also be expressed during only a specific stage of
25 development or life cycle of the host cell or host organism, again for example by expression under the control of a promoter that is specific for said stage of development or life cycle. Also, as already mentioned above, said expression may be constitutive, transient and/or inducible.

According to one specific embodiment, the expression of a nucleotide sequence of
30 the invention in a host cell or host organism may be partly or totally reduced (i.e. knocked out), compared to the original (e.g. native) host cell or host organism. This may for

instance be achieved in a transient manner using antisense and/or RNA-interference techniques well known in the art, or in a constitutive manner using random, site specific and/or chemical mutagenesis of the nucleotide sequence of the invention, or any other suitable techniques for generating "knock-down" or "knock-out" animals.

5 Suitable transformation techniques will be clear to the skilled person and may depend on the intended host cell/host organism and the genetic construct to be used. Some preferred, but non-limiting examples of suitable techniques include ballistic transformation, (micro-)injection, transfection (e.g. using suitable transposons), electroporation and lipofection. For these and other suitable techniques, reference is again
10 made to the handbooks and patent applications mentioned above.

 After transformation, a step for detecting and selecting those host cells or host organisms that have been successfully transformed with the nucleotide sequence/genetic construct of the invention may be performed. This may for instance be a selection step based on a selectable marker present in the genetic construct of the invention or a step
15 involving the detection of the amino acid sequence of the invention, e.g. using specific antibodies.

 The transformed host cell (which may be in the form of a stable cell line) or host organisms (which may be in the form of a stable mutant line or strain) form further aspects of the present invention.

20 Preferably, these host cells or host organisms are such that they express, or are (at least) capable of expressing (e.g. under suitable conditions), an amino acid sequence of the invention (and in case of a host organism: in at least one cell, part, tissue or organ thereof). The invention also includes further generations, progeny and/or offspring of the host cell or host organism of the invention, that may for instance be obtained by cell
25 division or by sexual or asexual reproduction.

 To produce/obtain expression of the amino acid sequences of the invention, the transformed host cell or transformed host organism may generally be kept, maintained and/or cultured under conditions such that the (desired) amino acid sequence of the invention is expressed/produced. Suitable conditions will be clear to the skilled person
30 and will usually depend upon the host cell/host organism used, as well as on the regulatory elements that control the expression of the (relevant) nucleotide sequence of

the invention. Again, reference is made to the handbooks and patent applications mentioned above in the paragraphs on the genetic constructs of the invention.

Generally, suitable conditions may include the use of a suitable medium, the presence of a suitable source of food and/or suitable nutrients, the use of a suitable temperature, and optionally the presence of a suitable inducing factor or compound (e.g. when the nucleotide sequences of the invention are under the control of an inducible promoter); all of which may be selected by the skilled person. Again, under such conditions, the amino acid sequences of the invention may be expressed in a constitutive manner, in a transient manner, or only when suitably induced.

It will also be clear to the skilled person that the amino acid sequence of the invention may (first) be generated in an immature form (as mentioned above), which may then be subjected to post-translational modification, depending on the host cell/host organism used. Also, the amino acid sequence of the invention may be glycosylated, again depending on the host cell/host organism used.

The amino acid sequence of the invention may then be isolated from the host cell/host organism and/or from the medium in which said host cell or host organism was cultivated, using protein isolation and/or purification techniques known per se, such as (preparative) chromatography and/or electrophoresis techniques, differential precipitation techniques, affinity techniques (e.g. using a specific, cleavable amino acid sequence fused with the amino acid sequence of the invention) and/or preparative immunological techniques (i.e. using antibodies against the amino acid sequence to be isolated).

In one embodiment, the amino acid sequence thus obtained may also be used to generate antibodies specifically against said sequence or an antigenic part or epitope thereof.

Such antibodies, which form a further aspect of the invention, may be generated in a manner known per se, for example as described in GB-A-2 357 768, US-A-5,693,492, WO 95/32734, WO 96/23882, WO 98/02456, WO 98/41633 and/or WO 98/49306, and/or as described in the prior art referred to above. Often, but not exclusively, such methods will involve immunizing an immunocompetent host with the pertinent amino acid sequence of the invention or an immunogenic part thereof (such as a specific epitope), in amount(s) and according to a regimen such that antibodies against said amino

acid sequence are raised, and then harvesting the antibodies thus generated, e.g. from blood or serum derived from said host.

For instance, polyclonal antibodies can be obtained by immunizing a suitable host such as a goat, rabbit, sheep, rat, pig or mouse with (an epitope of) an amino acid sequence of the invention, optionally with the use of an immunogenic carrier (such as bovine serum albumin or keyhole limpet hemocyanin) and/or an adjuvant such as Freund's, saponin, ISCOM's, aluminium hydroxide or a similar mineral gel, or keyhole limpet hemocyanin or a similar surface active substance. After a suitable immune response has been raised (usually within 1-7 days), the antibodies can be isolated from blood or serum taken from the immunized animal in a manner known per se, which optionally may involve a step of screening for an antibody with desired properties (i.e. specificity) using known immunoassay techniques, for which reference is again made to for instance WO 96/23882.

Monoclonal antibodies may, for example, be produced using continuous cell lines in culture, including hybridoma-based and similar techniques, again essentially as described in the above cited references. Accordingly, cells and cell lines that produce monoclonal antibodies against an amino acid sequence of the invention form a further aspect of the invention, as do methods for producing antibodies against amino acid sequences of the invention, which methods may generally involve cultivating such a cell and isolating the antibodies from the culture (medium), again using techniques known per se.

Also, Fab-fragments against the amino acid sequences of the invention (such as F(ab)₂, Fab' and Fab fragments) may be obtained by digestion of an antibody with pepsin or another protease, reducing disulfide-linkages and treatment with papain and a reducing agent, respectively. Fab-expression libraries may, for instance, be obtained by the method of Huse et al., 1989, Science 245:1275-1281.

In another embodiment, the nucleotide sequences of the invention, the amino acid sequences of the invention, and/or a host cell or host organism that expresses such an amino acid sequence, may also be used in an assay or assay method generally (including but not limited to diagnostic assays and/or assays to determining the presence and/or absence of specific mutations and/or genetic markers, for example to determine susceptibility for a condition or disease associated with such a mutation or marker), and in particular in an assay to identify and/or (further) develop compounds and/or other

factors that can modulate the (biological) activity of, and/or that can otherwise interact with, the amino acid sequences of the invention, and such uses form further aspects of the invention. As will be clear to the skilled person, in this context, the amino acid sequence of the invention will serve as a target for interaction with such a compound or factor

5 In this context, the terms “*modulate*”, “*modulation*”, “*modulator*” and “*target*” will have their usual meaning in the art, for which reference is *inter alia* made to the definitions given in WO 98/06737. Generally, a modulator is a compound or factor that can enhance, inhibit/reduce or otherwise alter, influence or affect (collectively referred to as “*modulation*”) a functional property of a biological activity or process (for example, 10 the biological activity of an amino acid sequence of the invention).

 In this context, the amino acid sequence of the invention may serve as a target for modulation *in vitro* (e.g. as part of an assay or screen) and/or for modulation *in vivo* (e.g. for modulation by a compound or factor that is known to modulate the target, which compound or factor may for example be used as an active compound for agrochemical, 15 veterinary and/or pharmaceutical use).

 For example, the amino acid sequences, host cells and/or host organisms of the invention may be used as part of an assay or screen that may be used to identify and/or develop modulators of the amino acid sequence of the invention, such as a primary screen (e.g. a screen used to identify modulators of the target from a set or library of test 20 chemicals with unknown activity with respect to the target) and/or a secondary assay (e.g. an assay used for validating hits from a primary screen and/or used in optimizing hit molecules, e.g. as part of hits-to-leads chemistry).

 For instance, such an assay or screen may be configured as an *in vitro* assay or screen, which will generally involve binding of the compound or factor to be tested as a 25 potential modulator for the target (hereinbelow also referred to as “test chemical”) to the target, upon which a signal generated by said binding is measured. Suitable techniques for such *in vitro* screening will be clear to the skilled person, and are for example described in Eldefrawi et al., (1987). FASEB J., Vol.1, pages 262-271 and Rauh et al., (1990), Trends in Pharmacol. Sci., vol.11, pages 325-329. For example, such an assay or 30 screen may be configured as a binding assay or screen, in which the test chemical is used to displace a detectable ligand from the target (e.g. a radioactive or fluorescent ligand),

upon which the amount of ligand displaced from the target by the modulator is determined.

Assays aimed at identification of small molecule inhibitors of protein kinases are well known. There is a large number of technologies that can be classified in two main approaches:

1. Detection based on an antibody specifically recognizing the phosphorylated product (peptide) of the kinase action

These approaches rely on the specificity of antibody to differentiate between the peptide containing a phosphorylated amino acid vs. the non-phosphorylated peptide.

This recognition can utilize, for example, the following readouts:

- fluorescence polarization (FP), where the detection is based on the change of the overall size upon antibody binding to fluorescent labeled peptide.
- fluorescence intensity (FI), where the antibody binding causes change in the intensity of the emission from a fluorescently labeled peptide.
- time-resolve fluorescence (where both the antibody and the peptide are labeled with different fluorescent labels and the binding leads to energy transfer between these two labels. This in turn leads to shortening of the donor label fluorescence lifetime. These assays have at least two different embodiments: biochemical or cell-based, via enzyme complementation (the latter is marketed as HitHunter (TM), by Discoverex) .

2. Detection based on other reagents specifically recognizing phosphate groups.

Recently, at least three new techniques have been developed that do not rely on the use of specific antibody.

- Trivalent metal ion containing beads. These bind specifically to phosphate groups. The detection is then performed by change in fluorescence polarization, the same as above, upon antibody binding. (IMAP(TM), Molecular Devices))
- based on chemical modification of the phosphate group and a specific label in the substrate to yield a change in fluorescence intensity. (IQ(TM), Pierce)
- based on chemical modification of the phosphate group to yield a biotin derivative. The detection is performed via FP upon avidin or streptavidin binding. (Caliper)"

Such an assay or screen may also be configured as a cell-based assay or screen, in which a host cell of the invention is contacted with/exposed to a test chemical, upon which at least one biological response by the host cell is measured.

Suitable cells or cell lines for such cell based assays include those mentioned
5 above. In one preferred, but non-limiting embodiment, the cell or cell line may be a mammalian, and in particular human, cell or cell line which is related to metabolic processes or metabolic disease and/or used as a cellular model for metabolic disease, including but not limited to liver cells or cell lines, adipocytes or muscle cells or cell lines such as HEPG2 cells, 3T3L1 adipocytes, CTC12 cells and L6 myotubes.

Also, such an assay or screen may also be configured as a whole animal screen,
10 in which a host organism of the invention is contacted with/exposed to a test chemical, upon which at least one biological response (such as a phenotypical, behavioural and/or physiological change, including but not limited to paralysis or death) by the host organism is measured. Such screens may be carried out in any model organism known
15 per se, including but not limited to yeast, *Drosophila*, zebrafish or *C. elegans*.

Thus, generally, the assays and screens described above will comprise at least one step in which the test chemical is contacted with the target (and/or with a host cell or host organism that expresses the target), and in particular in such a way that a signal is generated that is representative for the modulation of the target by the test chemical
20 (including modulation of the interaction of the target protein with any other proteins. Suitable techniques for screening such protein-protein interactions, such as FRET, SPA and BRET, will be clear to the skilled person). In a further step, said signal may then be detected.

Accordingly, in one aspect, the invention relates to a method for generating a
25 signal that is representative for the interaction of an amino acid sequence of the invention with a test chemical, said method at least comprising the steps of:

- a) contacting the amino acid sequence of the invention, or a host cell or host organism containing/expressing an amino acid sequence of the invention, with said test chemical, in such a way that a signal may be generated that is representative for the
30 interaction between said test chemical and said amino acid sequence; and optionally
- b) detecting the signal that may thus be generated.

In another aspect, the invention relates to a method for identifying modulators of an amino acid sequence of the invention (e.g. from a set or library of test chemicals), said method at least comprising the steps of:

- a) contacting the amino acid sequence of the invention, or a host cell or host organism containing/expressing an amino acid sequence of the invention, with a test chemical, in such a way that a signal may be generated that is representative for the interaction between said test chemical and said amino acid sequence; and optionally
- b) detecting the signal that may thus be generated, said signal identifying a modulator of said amino acid sequence.

Compounds that may be tested using the methods of the invention are generally described below.

The assays and screens of the invention may be carried out at medium throughput to high throughput, for example in an automated fashion using suitable robotics. In particular, in this embodiment, the method of the invention may be carried out by contacting the target with the test compound in a well of a multi-well plate, such as a standard 24, 96, 384, 1536 or 3456 well plate.

Usually, in a screen or assay of the invention, for each measurement, the target or host cell/host organism will be contacted with only a single test compound. However, it is also within the scope of the invention to contact the target with two or more test compounds - either simultaneously or sequentially - for example to determine whether said combination provides a synergistic effect.

Once a test chemical has been identified as a modulator for an amino acid sequence of the invention (e.g. by means of a screen or assay as described hereinabove), it may be used per se as a modulator of the amino relevant amino acid sequence of the invention (e.g. as an active substance for pharmaceutical use), or it may optionally be further optimized for final use, e.g. to improve properties such as solubility, ADME-TOX (see for example <http://www.scherago.com/admet/> or <http://www.lifesciencesinfo.com/admet>) and other desired properties. It will be clear to the skilled person that the nucleotide sequences, amino acid sequences, host cells/host organisms and/or methods of the invention may find further use in such optimization methodology, for example as (part of) secondary assays.

The invention is not particularly limited to any specific manner or mechanism in/via which the modulator (e.g. the test chemical, compound and/or factor) modulates, or interacts with, the target (*in vivo* and/or *in vitro*). For example, the modulator may comprise an agonist, an antagonist, an inverse agonist, a partial agonist, a competitive inhibitor, a non-competitive inhibitor, a cofactor, an allosteric inhibitor or other allosteric factor for the target, and/or may be a compound or factor that enhances or reduces binding of target to another biological component associated with its (biological) activity, such as another protein or polypeptide, a receptor, or a part or organelle of a cell. As such, the modulator may bind with the target (at the active site, at an allosteric site, at a binding domain and/or at another site on the target, e.g. covalently or via hydrogen bonding), block the active site of the target (in a reversible, irreversible or competitive manner), block a binding domain of the target (in a reversible, irreversible or competitive manner), and/or influence or change the conformation of the target.

As such, the test chemical/modulator may for instance be:

- an analog of a known substrate of the target;
- an oligopeptide, e.g. comprising between 2 and 20, preferably between 3 and 15 amino acid residues;
- an antisense or double stranded RNA molecule;
- a protein, polypeptide;
- a cofactor or an analog of a cofactor.

Preferably, the compound is an inhibitor of the target, although the invention in its broadest sense is not limited thereto.

The test chemical/modulator may also be a reference compound or factor, which may be a compound that is known to modulate or otherwise interact with the target (e.g. a known substrate or inhibitor for the target) or a compound or factor that is generally known to modulate or otherwise interact with other members from the general class to which the target belongs (e.g. a known substrate or inhibitor of said class).

Preferably, however, the compound(s) will be "small molecules", by which is generally meant herein a molecular entity with a molecular weight of less than 1500, preferably less than 1000. This may for example be an organic, inorganic or organometallic molecule, which may also be in the form of a suitable salt, such as a

water-soluble salt; and may also be a complex, chelate and/or a similar molecular entity, as long as its (overall) molecular weight is within the range indicated above.

In a preferred embodiment, such a "small molecule" has been designed according, and/or meets the criteria of, at least one, preferably at least any two, more preferably at least any three, and up to all of the so-called Lipinski rules for drug likeness prediction (vide Lipinski et al., *Advanced Drug Delivery Reviews* 23 (1997), pages 3-25). As is known in the art, small molecules which meet these criteria are particularly suited (as starting points) for the design and/or development of pharmaceuticals for human use, and may for instance be used as starting points for hits-to-leads chemistry, and/or as starting points for lead development (in which the methods of the invention may also be applied).

Also, for these purposes, the design of such small molecules (as well as the design of libraries consisting of such small molecules) will preferably also take into account the presence of pharmacophore points, for example according to the methods described by I. Muegge et al., *J. Med. Chem.* 44, 12 (2001), pages 1-6 and the documents cited herein.

The term "small peptide" generally covers (oligo)peptides that contain a total of between 2 and 35, such as for example between 3 and 25, amino acids (e.g. in one or more connected chains, and preferably a single chain). It will be clear that some of these small peptides will also be included in the term small molecule as used herein, depending on their molecular weight.

In one preferred, but non-limiting embodiment, the invention is used to screen a set or library of (related or otherwise unrelated) small molecules, for example a standard "robustness set", a primary screening library (e.g. of otherwise unrelated compounds), a combinatorial library, a series of closely related chemical analogos. Such sets or libraries will be clear to the skilled person, and may for instance include, but are not limited to, such commercially available chemical libraries such as the various libraries available from Tocris Cookson, Bristol, UK.

In yet a further aspect, the invention relates to a method for identifying a compound that can be used in (the preparation of a pharmaceutical composition for) the prevention and/or treatment of metabolic diseases (e.g. from a set or library of test chemicals), said method at least comprising the steps of:

- a) contacting an amino acid sequence of the invention, and/or a host cell or host organism containing/expressing an amino acid sequence of the invention, with a test chemical, in such a way that a signal may be generated that is representative for the interaction between said test chemical and said amino acid sequence of the invention;
5 and optionally
- b) detecting the signal that may thus be generated, said signal identifying a modulator of said amino acid sequence.

The modulators thus identified can be used in (the preparation of a pharmaceutical composition for) the prevention and/or treatment of metabolic diseases, and/or can be
10 used to develop other compounds that can be used in (the preparation of a pharmaceutical composition for) the prevention and/or treatment of metabolic diseases, i.e. as already outlined above.

The invention also relates to the use of an amino acid sequence and/or a host cell/host organism of the invention in such a method.

15 Also, as already mentioned above, the use of the human nucleotide sequences of SEQ ID NOS: 3, 5, 7 and/or 9 and/or of sequences derived therefrom (such as mutants, parts, fragments and/or fusions thereof as described hereinabove), of the human amino acid sequences of SEQ ID NOS: 4, 6, 8 and/or 10 and/or of sequences derived therefrom (such as analogs, parts, fragments, and/or fusions thereof as described hereinabove), and
20 of host cells/host organisms containing/expressing these, are usually preferred, in particular when the invention is used to develop compounds for pharmaceutical use.

As already mentioned above, the compounds and/or factors that have been identified and/or developed as modulators of the amino acid sequences of the invention (and/or precursors for such compounds) may be useful as active substances in the
25 pharmaceutical field, for example in the preparation of pharmaceutical compositions, and both such modulators as well as (pharmaceutical) compositions containing them comprise further aspects of the invention.

In particular, the compounds and composition of the invention may be used in (the preparation of pharmaceutical compositions for) the prevention (e.g. prophylaxis)
30 and/or treatment of metabolic diseases (which for the purposes herein in its broadest

sense also includes preventing, treating and/or alleviating the symptoms and/or complications of such metabolic diseases).

In particular, such compounds and composition may be used in (the preparation of pharmaceutical compositions for) the prevention (e.g. prophylaxis) and/or treatment of metabolic diseases (which for the purposes herein in its broadest sense also includes preventing, treating and/or alleviating the symptoms and/or complications of such metabolic diseases).

In particular, the compounds and compositions of the invention may be used for preventing and/or treating:

- hyperglycemic conditions and/or other conditions and/or diseases that are (primarily) associated with (the response or sensitivity to) insulin, including but not limited to all forms of diabetes and disorders resulting from insulin resistance, such as Type I and Type II diabetes, as well as severe insulin resistance, hyperinsulinemia, and hyperlipidemia, e.g., obese subjects, and insulin-resistant diabetes, such as Mendenhall's Syndrome, Werner Syndrome, leprechaunism, lipotrophic diabetes, and other lipotrophies;
- conditions caused or usually associated with hyperglycemic conditions and/or obesity, such as hypertension, osteoporosis and/or lipodystrophy.
- so-called "metabolic syndrome" (also known as "Syndrome X") which is a condition where several of the following conditions coexist: hypertension; insulin resistance; diabetes; dyslipidemia; and/or obesity.

In particular, the compounds and compositions of the invention may be used for preventing and/or treating diabetes, especially Type I and Type II diabetes. "Diabetes" itself refers to a progressive disease of carbohydrate metabolism involving inadequate production or utilization of insulin and is characterized by hyperglycemia and glycosuria.

Also, as mentioned above, the amino acid sequences of the invention and in particular the nucleotide sequences of the invention, and more in particular the human amino acid sequences and nucleotide sequences of the invention may be used for diagnostic purposes, for example as part of diagnostic assays and/or as part of kits for performing such assays (in which such a kit will comprise at least a nucleotide sequence of the invention) may be suitably packaged (e.g. in a suitable container) and may

optionally further comprise one or more elements for such kits known per se, such as suitable reagents, buffers or other solvents, and instructions for use).

In particular, the amino acid sequences and nucleotide sequences of the invention, as well as assays and kits using such sequences, may be used for diagnostic purposes relating to one or more of the metabolic diseases specified, for example as assays to determine the presence and/or absence in an individual of specific mutations and/or genetic markers that relate to one or more of the metabolic diseases specified above, to determine the susceptibility and/or any predisposition to any of said metabolic diseases in an individual, to determine if any genetically determined factors contribute or even cause (in full or in part) said metabolic disease in an individual, determine and/or to confirm the kind of metabolic disease from which an individual suffers, and/or to predict the further progress of such a metabolic disease in an individual. It will also be clear that any results obtained using such a diagnostic method or assay may also provide guidance to the clinician as to how such a metabolic disease should be treated in an individual, e.g. which medication should be prescribed and/or the dosage regimen to be used.

It should also be noted that, for the treatment of metabolic disease in humans, the compound used will usually and preferably be an inhibitor of an amino acid sequence of the invention, although the invention in its broadest sense is not limited thereto.

In one specific, but non-limiting, embodiment of the invention, a compound is considered an inhibitor of one of the amino acid sequences of the invention if, in a relevant assay such as the kinase activity assays referred to above (or a suitable modification thereof, for example using partially or fully purified protein), said compound reduces the activity of said amino acid sequence, i.e. by at least 1%, preferably at least 10%, such as by 20% or more, compared to the activity without the presence of said compound.

In an even more specific, but non-limiting, embodiment of the invention, a compound is considered an inhibitor of one of the amino acid sequences of the invention if, in a relevant assay, such as a binding assay, said compound has an IC₅₀ value of less than 1000 μ M, preferably at than 500 μ M, more preferably less than 250 μ M, even more preferably less than 100 μ M, for example 50 μ M or less, such as about 10 μ M or less.

Again, preferably, in the invention compounds are used that are modulators, and in particular inhibitors, of the human amino acid sequences of SEQ ID NO: 4, 6, 8 and/or 10, and/or of amino acid sequences derived therefrom, such as analogs, mutants, parts, fragments and/or fusions as described above.

5 For pharmaceutical use, the compounds of the invention may be used as a free acid or base, and/or in the form of a pharmaceutically acceptable acid-addition and/or base-addition salt (e.g. obtained with non-toxic organic or inorganic acid or base), in the form of a hydrate, solvate and/or complex, and/or in the form of a pre-drug, such as an ester. Such salts, hydrates, solvates, etc. and the preparation thereof will be clear to the skilled person; reference is for instance made to the salts, hydrates, solvates, etc.
10 described in US-A-6,372,778, US-A-6,369,086 and US-6,369,067

Generally, for pharmaceutical use, the compounds of the inventions may be formulated as a pharmaceutical preparation comprising at least one compound of the invention and at least one pharmaceutically acceptable carrier, diluent or excipient and/or
15 adjuvant, and optionally one or more further pharmaceutically active compounds. By means of non-limiting examples, such a formulation may be in a form suitable for oral administration, for parenteral administration (such as by intravenous, intramuscular or subcutaneous injection or intravenous infusion), for topical administration, for administration by inhalation, by a skin patch, by an implant, by a suppository, etc. Such
20 suitable administration forms - which may be solid, semi-solid or liquid, depending on the manner of administration - as well as methods and carriers for use in the preparation thereof, will be clear to the skilled person; reference is again made to for instance US-A-6,372,778, US-A-3,696, 086 and US-6,369,067.

The pharmaceutical preparations of the invention are preferably in a unit dosage
25 form, and may be suitably packaged, for example in a box, blister, vial, bottle, sachet, ampoule or in any other suitable holder or container (which may be properly labeled); optionally with one or more leaflets containing product information and/or instructions for use. Generally, such unit dosages will contain between 1 and 500 mg of the at least one compound of the invention, e.g. about 10, 25, 50, 100, 200, 500 or 1000 mg per unit
30 dosage.

For pharmaceutical use, at least one compound of the invention will generally be administered in an amount of between 0.01 to 150 mg/kg body weight per day of the patient, divided over one or more daily doses. The amount(s) to be administered and the further treatment regimen may be determined by the treating clinician, depending on factors such as the age, gender and general condition of the patient and the nature and severity of the disease/symptoms to be treated.

Thus, in a further aspect, the invention relates to a composition, and in particular a composition for pharmaceutical use, that contains at least one compound of the invention (i.e. a compound that has been identified, discovered and/or developed using a nematode or method as described herein) and at least one suitable carrier (i.e. a carrier suitable for pharmaceutical use). The invention also relates to the use of a compound of the invention in the preparation of such a composition.

Preferably, the compounds and compositions of the invention are administered orally and/or in a form intended and/or suitable for oral administration.

It is also envisaged that the above compounds and compositions may be of value in the veterinary field, which for the purposes herein not only includes the prevention and/or treatment of diseases in animals, but also - for economically important animals such as cattle, pigs, sheep, chicken, fish, etc. - enhancing the growth and/or weight of the animal and/or the amount and/or the quality of the meat or other products obtained from the animal. Thus, in a further aspect, the invention relates to a composition for veterinary use that contains at least one compound of the invention (i.e. a compound that has been identified, discovered and/or developed using a nematode or method as described herein) and at least one suitable carrier (i.e. a carrier suitable for veterinary use). The invention also relates to the use of a compound of the invention in the preparation of such a composition.

In the agrochemical field, the invention may be used to identify compounds suitable for use in pesticides, insecticides, nematicides and/or other biocides or plant protection agents. For example, the compounds of the invention may be used to control the species listed in US-A-6,372,774. For this purpose, the compounds of the invention (or a suitable salt, hydrate or ester thereof) may be suitably formulated with one or more agrochemically acceptable carriers, to provide a formulation suitable for agrochemical

use, as will be clear to the skilled person (reference is, for example, made to the formulations and uses described in US-A-6,372,774).

Thus, in a further aspect, the invention relates to a composition for agrochemical use that contains at least one compound of the invention (i.e. a compound that has been
5 identified, discovered and/or developed using a nematode or method as described herein) and at least one suitable carrier (i.e. a carrier suitable for agrochemical use). The invention also relates to the use of a compound of the invention in the preparation of such a composition.

The invention will now be further illustrated by means of the following non-
10 limiting Experimental Part.

In the Figures:

- Figure 1 schematically shows vector pGN49A (see also also WO 00/01846 and British patent application 0012233, both by Applicant);
- Figures 2A and 2B are photographs (enhanced using the Scion Image (Scion Corp)
15 software package) showing reduced fat-absorption phenotype in *C. elegans* upon Nile Red Staining: Figure 2A = reduced fat storage (T17E9.1a- invention); Figure 2B = reference (vector gGN29 without T17E9.1a).

Experimental part:

20 In the Experimental Part below, unless indicated otherwise, all steps for handling and cultivating *C. elegans* were performed using standard techniques and procedures, for which reference is made to the standard *C. elegans* handbooks, such as W.B. Wood et al., "*The nematode Caenorhabditis elegans*", Cold Spring Harbor Laboratory Press (1988); D.L. Riddle et al., "*C. ELEGANS II*", Cold Spring Harbor Laboratory Press (1997);
25 "*Caenorhabditis elegans, Modern Biological analysis of an organism*": ed. by H. Epstein and D. Shakes, Methods in Cell Biology, Vol 48, 1995; and "*C. elegans, a practical approach*", ed. by I.A. Hope, Oxford University Press Inc. New York, USA, 1999.

Downregulation of the gene(s) of interest in *C. elegans* was achieved by RNAi feeding techniques using an *E.coli* strain capable of expressing a dsRNA corresponding
30 to the gene(s) of interest, as generally described in - *inter alia* - the International application WO 00/01846 by applicant and the handbooks referred to above.

Also, unless indicated otherwise, all cloning and other molecular biology steps were performed using standard techniques and protocols, i.e. as provided by the manufacturers of the reagents/kits used and/or as described in the standard handbooks, such as Sambrook et al, "Molecular Cloning: A Laboratory Manual" (2nd.ed.), Vols. 1-3, Cold Spring Harbor Laboratory Press (1989) and F. Ausubel et al, eds., "Current protocols in molecular biology", Green Publishing and Wiley Interscience, New York (1987).

Fat accumulation in *C. elegans daf-2 (e1370)* was determined visually under a microscope upon staining with Nile-red, using an adaptation of the general methodology described by Ogg et al., Nature, Vol. 389, 994 (1997). For the general methodology, reference is also made to Thaden et al., 1999 International Worm Meeting abstract 837; Ashrafi and Ruvkun, 2000 East Coast Worm Meeting abstract 67; Ashrafi, Chang and Ruvkun, 2001 International Worm Meeting abstract 325; and Rottiers and Antebi, 2001 International Worm Meeting abstract 620 (all abstracts available from Worm Literature Index at <http://elegans.swmed.edu/wli/>).

Example 1: Preparation *E.coli* RNA feeding strain for expression of T17E9.1a double stranded RNA.

A vector for expression of dsRNA for downregulation of *C. elegans* gene T17E9.1a (*kin-18*) was prepared as follows.

The DNA fragment of SEQ ID NO:11, which corresponds to 1576 nucleotides (26 % compared to the genomic sequence) of the *C. elegans* T17E9.1a gene, was obtained by PCR from genomic *C. elegans* DNA, using the following primers:

- forward primer : CGAAAACCAGCAGAAGAGCG [SEQ ID NO: 12]
- reverse primer : TCGAGGCATGTTCTAGACTTCG [SEQ ID NO: 13]

This fragment was inserted in the *SrfI*-site of expression vector pGN49a pGN49A (Figure 1, see also WO 00/01846 and British patent application 0012233, both by Applicant). This vector contains two T7 promoters flanking the *SrfI*-site, allowing

transcription of a nucleotide sequence inserted into said *SrT*-site into double stranded RNA, upon binding of a T7 polymerase to said promoter (vide WO 00/01846).

The resulting vector, designated pGN49A-T17E9.1a, was transformed overnight into *E. coli* strain AB 309-105 (see EP-A-1 093 526 by applicant, page 17.).

5 To normalize the culture, 250 μ l of the overnight culture (1 ml) was transferred to a 96 well plate and the OD at 600 nm was measured (Fluostar Galaxy plate reader BMG), the remaining 750 μ l centrifuged down. Next the pellet was re-suspended in S-complete fed (S-complete supplemented with 0.1mg/ml ampiciline and 1 mM IPTG) and volume adjusted to obtain OD₆₀₀ value of 1

10 Example 2: Generation of fat storage phenotype in *C. elegans*.

In this example, *C. elegans* strain CB1370 containing the temperature sensitive daf-2 allele e-1370 is used (Ogg et al., supra). CB 1370 is publicly available from, for example, the Caenorhabditis Genetics Center (CGC), Minnesota, USA).

15 To generate the fat-storage phenotype, L1 worms of strain CB 1370 were cultivated at a temperature of 15 °C in S-Complete fed-medium in the wells of a 96 well plate (1 L1 nematodes per well) under essentially synchronized conditions, until the nematodes reached the L2 stage.

Then, the temperature was increased to 20°C, and the worms were further
20 cultivated at said temperature until their F1 offspring reached the L4 stage (about 144 hrs). Due to the presence of the daf-2 allele e-1370, this raise in temperature from 15°C to 20°C causes the nematodes to accumulate fat, mainly in their intestinal and hypodermal tissue (vide Ogg et al. and Figures 2A and 2B).

The accumulation of fat (in the form of droplets) was made visible by means of
25 Nile Red staining: L4 animals were washed several times with M9 (supplemented with 0.1% PEG) to remove the remaining *E.coli*, and fixed with MeOH (fc. 33%). After fixation the nematodes were stained with nile red (fc 0.375 mM in 37.5% MeOH) for 4 hours. MeOH and excess dye was removed through several washes with M9 (supplemented with 0.1% PEG). The staining pattern was visualized under UV using a
30 500 nm long pass filter.

For testing the influence of the gene T17E9.1a on fat storage, during the steps described above, the worms were grown on 30 μ l of the normalized *E.coli* pGN49A-T17E9.1a strain of Example 1 ($OD_{600} = 1$) as a food source. As a reference, the daf-2 (e1370) nematodes were grown in a similar manner, but with *E. coli* strain AB 309-105 containing vector pGN29A without the T17E9.1a insert as a food source, used in the same amount. All samples were carried out in quadruplicate.

The results were as follows: worms fed on *E.coli* pGN49A-T17E9.1a strain, which downregulates the expression of T17E9.1a through RNA interference, showed a strong reduction of the accumulation of fat, compared to the reference (vide Figures 2A and 2B).

These results show that T17E9.1a is involved in the regulation of the (daf-2 dependent) *daf*-pathway and the accumulation of fat in the nematode. It is known in the art that both are models for insulin resistance and fat handling in mammals, such as humans.

Bio-informatic analysis of the sequence of T17E9.1a (kin-18) showed that the amino acid sequence of the kin-18 protein (SEQ ID NO:2) has the following degree of sequence identity (in percentages at the amino acid level) with the human amino acid sequences of SEQ ID NOS: 4, 6, 8 and 10, respectively:

- SEQ ID NO:4 ("JIK"): overall sequence: 36%; kinase domain: 61%
- SEQ ID NO:6 ("PSK"): overall sequence: 24%; kinase domain: 59%
- SEQ ID NO:8 ("TAOI"): overall sequence: 32%; kinase domain: 59%
- SEQ ID NO:10 ("Q9P2I6"): overall sequence: 33%; kinase domain: 61%

Example 3: Screening of SEQ ID NO. 4 ("JIK").

A protein with the amino acid sequence of SEQ ID NO: 4, isolated and purified in a manner known per se, were used to screen a reference set of 96 commercially available kinase inhibitors (obtained from several commercial sources such as Acros Organics, Belgium/USA; Alexis Biochemicals, UK; Biomol, USA; Calbiochem, USA; ICN Biomedicals; Sigma; and Tocris, UK), using the commercially available Kinase Glo® Luminescent Kinase Assay (Promega, USA), essentially according to the instructions of the manufacturer.

In this assay, 6 out of 96 compounds showed a percentage which differed more than 3 Standard Deviation (SD) from 0 (zero) inhibition. These 6 compounds were designated as hits, corresponding with a hit-rate of about 6 %.

Example 4: Screening of SEQ ID NO. 8 ("TAOI").

A protein with the amino acid sequence of SEQ ID NO: 8, isolated and purified in a manner known per se, were used to screen a reference set of 96 commercially available kinase inhibitors (obtained from several commercial sources such as Acros Organics, Belgium/USA; Alexis Biochemicals, UK; Biomol, USA; Calbiochem, USA; ICN Biomedicals; Sigma; and Tocris, UK), using the commercially available Kinase Glo® Luminescent Kinase Assay (Promega, USA), essentially according to the instructions of the manufacturer.

In this assay, 6 out of 96 compounds showed a percentage which differed more than 3 Standard Deviation (SD) from 0 (zero) inhibition. These 6 compounds were designated as hits, corresponding with a hit-rate of about 6 %.

CLAIMS:

- 5 1. Nucleic acid, in essentially isolated form, said nucleic acid encoding one of the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and/or SEQ ID NO: 10, or an analog, variant, allele, ortholog, part and/or fragment of one of the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and/or SEQ ID NO: 10.
- 10 2. Nucleic acid, in essentially isolated form, said nucleic acid comprising one of the nucleotide sequences of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7 and/or SEQ ID NO: 9, or a mutant, variant, allele, analog, ortholog, part and/or fragment thereof.
- 15 3. Genetic construct, comprising the nucleic acid of claim 1 and/or of claim 2, and optionally one or more further elements of genetic constructs known per se.
- 20 4. Host cell or host organism, transformed with and/or containing a nucleic acid according to claim 1 and/or claim 2 and/or a genetic construct according to claim 3.
- 25 5. Host cell or host organism, (capable of) expressing and/or producing one of the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and/or SEQ ID NO: 10 or an analog, variant, allele, analog, ortholog, part and/or fragment thereof.
- 30 6. Method for producing one of the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and/or SEQ ID NO: 10, or an analog, variant, allele, ortholog, part and/or fragment thereof, said method comprising at least the steps of:
- a) expressing of a nucleic acid according to claim 1 and/or 2 in a suitable host cell or host organism; and optionally

b) isolating the amino acid sequence thus expressed.

7. Method for producing one of the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and/or SEQ ID NO: 10, or an analog, variant, allele, ortholog, part and/or fragment thereof, said method comprising at least the steps of:

- 5 a) maintaining and/or cultivating a host cell or host organism according to claim 4 or 5 under conditions such that said host cell or host organism expresses or produces one of the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and/or SEQ ID NO: 10, or an analog, variant, allele, ortholog, part and/or fragment thereof; and optionally
- 10 b) isolating the amino acid sequence thus expressed/produced.

8. Protein or polypeptide, in essentially isolated form, said protein or polypeptide comprising one of the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and/or SEQ ID NO: 10, or an analog, variant, allele, ortholog, part and/or fragment thereof.

15

9. Method for generating a signal that is representative for the interaction of a protein or polypeptide of claim 8 with a test chemical, said method at least comprising the steps of:

20

- a) contacting a protein or polypeptide of claim 8, or a host cell or host organism of claim 4 or 5, with said test chemical, in such a way that a signal may be generated that is representative for the interaction between said test chemical and said protein or polypeptide; and optionally
- 25 b) detecting the signal that may thus be generated.

10. Method for identifying a modulator of a protein or polypeptide of claim 8, for example from a set or library of test chemicals, said method at least comprising the steps of:

30

- a) contacting a protein or polypeptide of claim 8, or a host cell or host organism of claim 4 or 5, with said test chemical, in such a way that a signal may be generated that is representative for the interaction between said test chemical and said protein or polypeptide; and optionally
- 5 b) detecting the signal that may thus be generated, said signal identifying a modulator of said amino acid sequence.

11. Modulator of a protein or polypeptide according to claim 8, identified and/or developed using a nucleic acid according to claim 1 and/or 2, a host cell according to
10 claim 4 and/or 5, a protein or polypeptide according to claim 8, and/or a method according to claim 9 and/or 10.

12. Modulator according to claim 11, being an inhibitor of a protein or polypeptide of claim 8.

15

13. Pharmaceutical composition, comprising at least one modulator according to claim 11 and/or 12 and at least one pharmaceutically acceptable carrier.

14. Pharmaceutical composition according to claim 13, being a composition
20 suitable and/or intended for oral administration.

15. Use of a modulator according to claim 11 and/or 12 in the preparation of a pharmaceutical composition.

25 16. Use of a modulator according to claim 11 and/or 12 in the preparation of a pharmaceutical composition for the prevention and/or treatment of metabolic diseases.

17. Antibody against a protein or polypeptide according to claim 8.



Figure 1

Figure 2A**Figure 2B**

SEQUENCE LISTING

5 <110> Devgen N.V.
 Devgen N.V.

10 <120> Kinase sequences useful for developing compounds for the
 prevention and/or treatment of metabolic diseases and nucleotide
 sequences encoding such kinase sequences.

 <130> P 02-011

15 <160> 13
 <170> PatentIn version 3.1

20 <210> 1
 <211> 2949
 <212> DNA

25 <213> Caenorhabditis elegans

 <400> 1
 atggcgccctg ccgtcttaca aaaacccggt gttatcaagg atccatcgat tgctgcattg 60
 30 ttcagtaata aggatccaga gcagagatat caagatttaa gagaaattgg acatggatct 120
 tttggagctg tctattttgc atatgacaaa aaaaatgagc agactgttgc gattaaaaag 180
 35 atgaatttta gtggaaaaca ggctgtcgaa aaatggaatg atattcttaa agaagtgtct 240
 tttctgaata cagttgttca tccacatatt gtcgactaca aggcttggtt tcttaaggac 300
 actacatggt ggcttgtgat ggagtactgt attggctctg cagccgatat agtggatgtc 360
 40 ttgcgaaaag gaatgcgaga agtcgaaatc gctgcgattt gctctcaaac tttggatgct 420
 cttcgatatc ttcactctct gaagcgaata catcgagata ttaaagctgg aaatattctg 480
 ctatctgata atgctattgt taaactagct gatttcggat ccgcatccct ggtagatccg 540
 45 gctcaaactt tcatcggaac gccgtttttc atggccccag aggtaattct ggcaatggat 600
 gagggtcact acacggatcg tgcagatatt tggtcattgg gtatcacgtg tatagagctg 660
 50 gccgaacgtc gtccaccatt gttcagtatg aatgcaatgt ctgccctcta ccatattgct 720
 caaaatgata ctccaactct ttctccaatt gacactagcg aacaaccgga atggctcgctg 780
 gaattcgttc aatttataga caaatgtctt cgaaaaccag cagaagagcg aatgtcagct 840
 55 gaagaatgct ttcgacatcc attcattcaa cgggtctcgcc catcagacac aattcaggaa 900

ctcatcaga gaacgaaaaa tatggtatta gagttggata attttcaata caaaaagatg 960
 agaaaactca tgtatttgga tgaaacagaa ggaaaagaag gaagtgaagg aaatggagca 1020
 5 tctgatgatt tagattttca tggaaatgaa gctaattcaa ttggaagagc aggagattct 1080
 gcgtcatctc gaagtgcttc tcttacttct ttccgatcaa tgcagagtag tggaggagct 1140
 10 ggtcttttag tgtccaccaa tacgacgggt gctatggata atgtgcatgg atcctctgga 1200
 tacggtaatg gaagtagttc gacgacgagc tccgcacgcc gccgtcctcc aattccttcg 1260
 caaatgctct cttctacatc aacgtctgggt gttggaacta tgccgagtca tggatcagtt 1320
 15 ggagcatcga ttacggcgat cgcagtcaat ccaacaccgt ctcttcaga acctatccca 1380
 acatcacaaac caacatcgaa atcagaatca tcttctatac tcgaaactgc acacgatgat 1440
 cctttggaca cgtcgatacg tgctccagtg aaagacttgc atatgccgca tcgagcagtc 1500
 20 aaggaacgaa tagccacgtt gcaaaatcac aaattcgca cgcttcgttc ccagagaata 1560
 atcaatcagg aacaagaaga atatacgaag gagaaacata tgtatgagca aatgagcaag 1620
 25 tacaagcatc tacgacaagc acatcacaaa gagctccaac aatttgaaga acgatgtgca 1680
 ttagatagag agcaactgag tgtgaaaatg gatcgagaac tcgaacaatt gacaacgaca 1740
 tactcgaaag aaaagatgag agtgaggtgt tcacagaata atgaactaga caaacggaaa 1800
 30 aaagatatcg aagatgggga gaaaaagatg aaaaagacga aaaatagtca aaatcagcag 1860
 cagatgaaac tgtattcagc gcaacaattg aaagaataca agtataacaa ggaggcacag 1920
 35 aaaacacgat tacgaagtct gaacatgcct cgaagtactt atgagaacgc aatgaaagaa 1980
 gtgaaagccg atctgaatcg agtgaaagat gcacgggaaa atgattttga cgagaagctt 2040
 cgtgcagaac ttgaagatga aattgtaagg tatcgaggc aacaactcag taatcttcac 2100
 40 caattggaag aacaattgga tgatgaagac gtaaacgtgc aagaacgcca aatggacacg 2160
 cgtcacggat tactgtcaaa gcagcatgaa atgacgcgag atttggaaat acagcatctc 2220
 45 aacgagcttc acgcgatgaa aaaacgacat ttggagacac aacacgaggc ggaatcggca 2280
 agtcaaaatg agtacacaca gaggcaacag gatgaattga gaaaaaagca tgcgatgcag 2340
 tcaagacaac agccaagaga tttaaagatc caagaagcac aaattcgaaa acaataccga 2400
 50 caagttgtga agactcagac tcgccaattt aagctctacc ttacacaaat ggtgcaagta 2460
 gttccaaaag atgaacaaaa agagctcacg tctcgactaa aacaggatca aatgcaaaaa 2520
 55 gtcgcacttc ttgcttcaca atacgaaagt caaatcaaaa aaatggttca ggataagaca 2580
 gtgaagctcg agtcgtggca agaagatgaa caacgggttc ttagtgagaa gttggagaaa 2640

gaattggaag aattgattgc ttatcagaag aagacgagag ccacattaga agagcagatt 2700
 5 aaaaaggaac gtacggcact cgaagaacga attggcacac gacgtgcaat gcttgaacag 2760
 aagattattg aagaacgcga acaaatggga gaaatgcgtc gactaaagaa ggagcaaatac 2820
 cgtgatcgac acagtcaaga acgccatcgt ctcgagaatc atttcgtacg gacgggctcg 2880
 10 acgagcagaa gttctggtgg gatcgctcct ggtggtggga attcaagcag tattcagatg 2940
 gctatgtag 2949

15 <210> 2

<211> 982

<212> PRT

20

<213> *Caenorhabditis elegans*

<400> 2

25 Met Ala Pro Ala Val Leu Gln Lys Pro Gly Val Ile Lys Asp Pro Ser
1 5 10 15

Ile Ala Ala Leu Phe Ser Asn Lys Asp Pro Glu Gln Arg Tyr Gln Asp
20 25 30

30

Leu Arg Glu Ile Gly His Gly Ser Phe Gly Ala Val Tyr Phe Ala Tyr
35 40 45

35 Asp Lys Lys Asn Glu Gln Thr Val Ala Ile Lys Lys Met Asn Phe Ser
50 55 60

Gly Lys Gln Ala Val Glu Lys Trp Asn Asp Ile Leu Lys Glu Val Ser
65 70 75 80

40 Phe Leu Asn Thr Val Val His Pro His Ile Val Asp Tyr Lys Ala Cys
85 90 95

Phe Leu Lys Asp Thr Thr Cys Trp Leu Val Met Glu Tyr Cys Ile Gly
100 105 110

45

Ser Ala Ala Asp Ile Val Asp Val Leu Arg Lys Gly Met Arg Glu Val
115 120 125

50 Glu Ile Ala Ala Ile Cys Ser Gln Thr Leu Asp Ala Leu Arg Tyr Leu
130 135 140

His Ser Leu Lys Arg Ile His Arg Asp Ile Lys Ala Gly Asn Ile Leu
145 150 155 160

55 Leu Ser Asp His Ala Ile Val Lys Leu Ala Asp Phe Gly Ser Ala Ser
165 170 175

Leu Val Asp Pro Ala Gln Thr Phe Ile Gly Thr Pro Phe Phe Met Ala
 180 185 190
 5 Pro Glu Val Ile Leu Ala Met Asp Glu Gly His Tyr Thr Asp Arg Ala
 195 200 205
 Asp Ile Trp Ser Leu Gly Ile Thr Cys Ile Glu Leu Ala Glu Arg Arg
 210 215 220
 10 Pro Pro Leu Phe Ser Met Asn Ala Met Ser Ala Leu Tyr His Ile Ala
 225 230 235 240
 Gln Asn Asp Pro Pro Thr Leu Ser Pro Ile Asp Thr Ser Glu Gln Pro
 245 250 255
 15 Glu Trp Ser Leu Glu Phe Val Gln Phe Ile Asp Lys Cys Leu Arg Lys
 260 265 270
 Pro Ala Glu Glu Arg Met Ser Ala Glu Glu Cys Phe Arg His Pro Phe
 275 280 285
 20 Ile Gln Arg Ser Arg Pro Ser Asp Thr Ile Gln Glu Leu Ile Gln Arg
 290 295 300
 25 Thr Lys Asn Met Val Leu Glu Leu Asp Asn Phe Gln Tyr Lys Lys Met
 305 310 315 320
 Arg Lys Leu Met Tyr Leu Asp Glu Thr Glu Gly Lys Glu Gly Ser Glu
 325 330 335
 30 Gly Asn Gly Ala Ser Asp Asp Leu Asp Phe His Gly Asn Glu Ala Asn
 340 345 350
 Ser Ile Gly Arg Ala Gly Asp Ser Ala Ser Ser Arg Ser Ala Ser Leu
 355 360 365
 Thr Ser Phe Arg Ser Met Gln Ser Ser Gly Gly Ala Gly Leu Leu Val
 370 375 380
 40 Ser Thr Asn Thr Thr Gly Ala Met Asp Asn Val His Gly Ser Ser Gly
 385 390 395 400
 Tyr Gly Asn Gly Ser Ser Ser Thr Thr Ser Ser Ala Arg Arg Arg Pro
 405 410 415
 45 Pro Ile Pro Ser Gln Met Leu Ser Ser Thr Ser Thr Ser Gly Val Gly
 420 425 430
 Thr Met Pro Ser His Gly Ser Val Gly Ala Ser Ile Thr Ala Ile Ala
 435 440 445
 Val Asn Pro Thr Pro Ser Pro Ser Glu Pro Ile Pro Thr Ser Gln Pro
 450 455 460
 55 Thr Ser Lys Ser Glu Ser Ser Ser Ile Leu Glu Thr Ala His Asp Asp
 465 470 475 480

Pro Leu Asp Thr Ser Ile Arg Ala Pro Val Lys Asp Leu His Met Pro
 .485 490 495
 5 His Arg Ala Val Lys Glu Arg Ile Ala Thr Leu Gln Asn His Lys Phe
 500 505 510
 Ala Thr Leu Arg Ser Gln Arg Ile Ile Asn Gln Glu Gln Glu Glu Tyr
 515 520 525
 10 Thr Lys Glu Asn Asn Met Tyr Glu Gln Met Ser Lys Tyr Lys His Leu
 530 535 540
 Arg Gln Ala His His Lys Glu Leu Gln Gln Phe Glu Glu Arg Cys Ala
 545 550 555 560
 15 Leu Asp Arg Glu Gln Leu Arg Val Lys Met Asp Arg Glu Leu Glu Gln
 565 570 575
 Leu Thr Thr Thr Tyr Ser Lys Glu Lys Met Arg Val Arg Cys Ser Gln
 580 585 590
 Asn Asn Glu Leu Asp Lys Arg Lys Lys Asp Ile Glu Asp Gly Glu Lys
 595 600 605
 25 Lys Met Lys Lys Thr Lys Asn Ser Gln Asn Gln Gln Gln Met Lys Leu
 610 615 620
 Tyr Ser Ala Gln Gln Leu Lys Glu Tyr Lys Tyr Asn Lys Glu Ala Gln
 625 630 635 640
 30 Lys Thr Arg Leu Arg Ser Leu Asn Met Pro Arg Ser Thr Tyr Glu Asn
 645 650 655
 Ala Met Lys Glu Val Lys Ala Asp Leu Asn Arg Val Lys Asp Ala Arg
 660 665 670
 35 Glu Asn Asp Phe Asp Glu Lys Leu Arg Ala Glu Leu Glu Asp Glu Ile
 675 680 685
 40 Val Arg Tyr Arg Arg Gln Gln Leu Ser Asn Leu His Gln Leu Glu Glu
 690 695 700
 Gln Leu Asp Asp Glu Asp Val Asn Val Gln Glu Arg Gln Met Asp Thr
 705 710 715 720
 45 Arg His Gly Leu Leu Ser Lys Gln His Glu Met Thr Arg Asp Leu Glu
 725 730 735
 Ile Gln His Leu Asn Glu Leu His Ala Met Lys Lys Arg His Leu Glu
 740 745 750
 50 Thr Gln His Glu Ala Glu Ser Ala Ser Gln Asn Glu Tyr Thr Gln Arg
 755 760 765
 55 Gln Gln Asp Glu Leu Arg Lys Lys His Ala Met Gln Ser Arg Gln Gln
 770 775 780

Pro Arg Asp Leu Lys Ile Gln Glu Ala Gln Ile Arg Lys Gln Tyr Arg
 785 790 795 800
 5 Gln Val Val Lys Thr Gln Thr Arg Gln Phe Lys Leu Tyr Leu Thr Gln
 805 810 815
 Met Val Gln Val Val Pro Lys Asp Glu Gln Lys Glu Leu Thr Ser Arg
 820 825 830
 10 Leu Lys Gln Asp Gln Met Gln Lys Val Ala Leu Leu Ala Ser Gln Tyr
 835 840 845
 Glu Ser Gln Ile Lys Lys Met Val Gln Asp Lys Thr Val Lys Leu Glu
 850 855 860
 15 Ser Trp Gln Glu Asp Glu Gln Arg Val Leu Ser Glu Lys Leu Glu Lys
 865 870 875 880
 Glu Leu Glu Glu Leu Ile Ala Tyr Gln Lys Lys Thr Arg Ala Thr Leu
 885 890 895
 20 Glu Glu Gln Ile Lys Lys Glu Arg Thr Ala Leu Glu Glu Arg Ile Gly
 900 905 910
 25 Thr Arg Arg Ala Met Leu Glu Gln Lys Ile Ile Glu Glu Arg Glu Gln
 915 920 925
 Met Gly Glu Met Arg Arg Leu Lys Lys Glu Gln Ile Arg Asp Arg His
 930 935 940
 30 Ser Gln Glu Arg His Arg Leu Glu Asn His Phe Val Arg Thr Gly Ser
 945 950 955 960
 Thr Ser Arg Ser Ser Gly Gly Ile Ala Pro Gly Val Gly Asn Ser Ser
 965 970 975
 35 Ser Ile Gln Met Ala Met
 980
 40 <210> 3
 <211> 4188
 45 <212> DNA
 <213> Homo sapiens
 <300>
 50 <308> genbank
 <309> 2002-10-04
 55 <313> (1)..(4188)

<300>
<308> genbank NM_016281
5 <309> 2002-10-04
<313> (1)..(4188)
<400> 3
10 gccggggaac aagccacagg agagcgactc aggaacaagt gtgggagagg aagcggcggc 60
ggcggcgccg ggcccggggg tgggtgacagc aggtctgagg ttgcatcata aatacaaagg 120
15 actgaagtta taaaagagaa aagagaagtt tgctgctaaa atgaatctga gcaatatgga 180
atatatttggt ccacacacaa aaagggtactg aagattttacc ccccaaaaaa aattgtcaat 240
20 gagaaataaa gctaactgat atcaaaaagc agagcctgct ctactggcca tcatgcgtaa 300
aggggtgctg aaggacccag agattgccga tctattctac aaagatgatc ctgaggaact 360
ttttattggt ttgcatgaaa ttggacatgg aagttttgga gcagtttatt ttgctacaaa 420
25 tgctcacacc aatgaggtgg tggcaattaa gaagatgtcc tatagtggga agcagaccca 480
tgagaaatgg caagatattc ttaaggaagt taaattttta cgacaattga agcatcctaa 540
tactattgag tacaagggt gttacttgaa agaacacact gcttggttgg tgatggaata 600
30 ttgcttaggc tcagcctctg atttattaga agttcataaa aaaccacttc aggaagtgga 660
gatcgctgcc attactcatg gagccttgca tggactagcc tacctacatt ctcatgcatt 720
35 gattcatagg gatattaaag caggaaatat tcttctaaca gagccaggtc aggtaaaact 780
agctgatttt ggatctgctt caatggcttc tcctgccaac tccttcgtgg gcacacctta 840
ctggatggct ccagaggtga tcttagctat ggatgaagga cagtatgatg ggaaagttga 900
40 tatttggtca cttggcatca cttgtattga attggcgga cggaagccgc cccttttcaa 960
catgaatgca atgagtgctt tatatcacat tgcccagaat gactcccaa cgttacagtc 1020
45 taatgaatgg acagactcct ttaggagatt tggtgattac tgcttgacaga aaatacctca 1080
ggaaaggcca acatcagcag aactattaag gcatgacttt gttcgacgag accggccact 1140
acgtgtcctc attgacctca tacagaggac aaaagatgca gttcgtgagc tagataacct 1200
50 acagtaccga aaaatgaaaa aaatactttt ccaagagaca cggaatggac ccttgaatga 1260
gtcacaggag gatgaggaag acagtgaaca tggaaaccagc ctgaacaggg aaatggacag 1320
55 cctgggcagc aaccattcca ttccaagcat gtccgtgagc acaggcagcc agagcagcag 1380
tgtgaacagc atgcaggaag tcatggacga gagcagttcc gaacttgtca tgatgcacga 1440

	tgacgaaagc	acaatcaatt	ccagctcctc	cgctcgtgcat	aagaaagatc	atgtattcat	1500
5	aagggatgag	gcgggcccacg	gcgatcccag	gcctgagccg	cggcctaccc	agtcagttca	1560
	gagccaggcc	ctccactacc	ggaacagaga	gcgctttgcc	acgatcaa	cagcatcttt	1620
	ggttacacga	cagatccatg	agcatgagca	ggagaacgag	ttgcgggaac	agatgtcagg	1680
10	ttataagcgg	atgcggcgcc	agcaccagaa	gcagctgata	gccctggaga	acaagctgaa	1740
	ggctgagatg	gacgagcacc	gcctcaagct	acagaaggag	gtggagacgc	atgccaaaca	1800
	ctcgtccatc	gagctggaga	agctggccaa	gaagcaagtg	gctatcatag	aaaaggaggc	1860
15	aaaggtagct	gcagcagatg	agaagaagtt	ccagcaacag	atcttggccc	agcagaagaa	1920
	agattttgaca	actttcttag	aaagtcagaa	gaagcagtat	aagattttgta	aggaaaaaat	1980
20	aaaagaggaa	atgaatgagg	accatagcac	acccaagaaa	gagaagcaag	agcggatctc	2040
	caaacataaa	gagaacttgc	agcacacaca	ggctgaagag	gaagcccacc	ttctcactca	2100
	acagagactg	tactacgaca	aaaattgtcg	tttcttcaag	cggaaaataa	tgatcaagcg	2160
25	gcacgaggtg	gagcagcaga	acattcggga	ggaactaaat	aaaaagagga	cccagaagga	2220
	gatggagcat	gccatgctaa	tccggcacga	cgagtccacc	cgagagctag	agtacaggca	2280
30	gctgcacacg	ttacagaagc	tacgcatgga	tctgatccgt	ttacagcacc	agacggaact	2340
	ggaaaaccag	ctggagtaca	ataagaggcg	agaaagagaa	ctgcacagaa	agcatgtcat	2400
	ggaacttcgg	caacagccaa	aaaacttaaa	ggccatggaa	atgcaaatta	aaaaacagtt	2460
35	tcaggacact	tgcaaagtac	agaccaaaca	gtataaagca	ctcaagaatc	accagttgga	2520
	agttactcca	aagaatgagc	acaaaacaat	cttaaagaca	ctgaaagatg	agcagacaag	2580
40	aaaacttgcc	attttggcag	agcagtatga	acagagtata	aatgaaatga	tggcctctca	2640
	agcgttacgg	ctagatgagg	ctcaagaagc	agaatgccag	gccttgaggc	tacagctcca	2700
	gcaggaaatg	gagctgctca	acgcctacca	gagcaaaatc	aagatgcaaa	cagaggcaca	2760
45	acatgaacgt	gagctccaga	agctagagca	gagagtgtct	ctgcgcagag	cacaccttga	2820
	gcagaagatt	gaagaggagc	tggctgccct	tcagaaggaa	cgcagcgaga	gaataaagaa	2880
50	cctattggaa	aggcaagagc	gagagattga	aacttttgac	atggagagcc	tcagaatggg	2940
	at ttgggaat	ttgggttacat	tagattttcc	taaggaggac	tacagatgag	attaaatttt	3000
	ttgccattta	caaaaaaaaa	aaaaaaaaaga	aaacagaaaa	aaattcagac	cctgcaaaac	3060
55	cacattcccc	attttaacgg	gcgttgctct	cactctctct	ctctcttact	cttactgaca	3120

tcgtgtcgga ctagtgctg tttattctta ctccatcagg ggcccccttc ctccccccgt 3180
gtcaactttc agtgctggcc aaaacctggc cgtctcttct attcacagta cacgtcacag 3240
5 tattgatgtg attcaaatg tttcagtga aactttggag acagttttaa caaaaccaat 3300
aaaccaacaa caaaaaaagt ggatgtatat tgctttaagc aatcactcat taccaccaat 3360
ctgtgaaagt aaagcaaaaa ataataataa taaatgccaa gggggagaga gacacaatat 3420
10 ccgcagcctt acaccttaac tagctgctgc attattttat tttattttat ttttttggtta 3480
tttattcatc aggaataaaa aaaacaaagt tttattaaag attgaaaatt tgatacatat 3540
15 tacagaaact aattgtgatg tacatatcag tggtgacata ttattacttt tttggggacg 3600
ggggtgggtg ggggtgaagag atcttgatg ttttagactgc tgcagagtta acttgctctca 3660
gcatactctga tgtatcataa tcatttctgc tgtgcagagg agggatacac ttaggggctc 3720
20 acagatccca gtagcacaat tgggctttgg caaatgggta ttttgtgtat agaggaattt 3780
aaggagaggt attacttatt ttcatatgtt attttaactg tttctcggat caaatttttt 3840
25 aacttcttct tcgtgttctt cccacctcc ttccttttcc agttcagtat ttggagttca 3900
acactgtctc tcaatcagat catctggatc tttttcttta tctcccttcc ccttcctaag 3960
tcccatttct tggtcataaa tattgcatta ttcacacttt caaactgtgt attttcttac 4020
30 aataaaaaat gatgaaaaaa aaaaaggctt tacttctttt gcatgcactt taaaaacaaa 4080
acaaaacatt tttcaggttc caaggaagag catgataact gtcagagctt ttaattatat 4140
35 ttgtaaataa aagtgttcat cacaaaaaaa aaaaaaaaaa aaaaaaaaaa 4188

<210> 4

40 <211> 898

<212> PRT

45 <213> Homo sapiens

<300>

50 <308> Genbank NP_057365.2

<309> 2002-10-01

<313> (1) .. (898)

55

<400> 4

	Met	Arg	Lys	Gly	Val	Leu	Lys	Asp	Pro	Glu	Ile	Ala	Asp	Leu	Phe	Tyr	
	1				5					10					15		
5	Lys	Asp	Asp	Pro	Glu	Glu	Leu	Phe	Ile	Gly	Leu	His	Glu	Ile	Gly	His	
				20					25					30			
	Gly	Ser	Phe	Gly	Ala	Val	Tyr	Phe	Ala	Thr	Asn	Ala	His	Thr	Asn	Glu	
			35					40					45				
10	Val	Val	Ala	Ile	Lys	Lys	Met	Ser	Tyr	Ser	Gly	Lys	Gln	Thr	His	Glu	
		50					55					60					
	Lys	Trp	Gln	Asp	Ile	Leu	Lys	Glu	Val	Lys	Phe	Leu	Arg	Gln	Leu	Lys	
	65					70					75					80	
15	His	Pro	Asn	Thr	Ile	Glu	Tyr	Lys	Gly	Cys	Tyr	Leu	Lys	Glu	His	Thr	
					85					90					95		
	Ala	Trp	Leu	Val	Met	Glu	Tyr	Cys	Leu	Gly	Ser	Ala	Ser	Asp	Leu	Leu	
20				100					105					110			
	Glu	Val	His	Lys	Lys	Pro	Leu	Gln	Glu	Val	Glu	Ile	Ala	Ala	Ile	Thr	
			115					120					125				
25	His	Gly	Ala	Leu	His	Gly	Leu	Ala	Tyr	Leu	His	Ser	His	Ala	Leu	Ile	
		130					135					140					
	His	Arg	Asp	Ile	Lys	Ala	Gly	Asn	Ile	Leu	Leu	Thr	Glu	Pro	Gly	Gln	
	145					150					155					160	
30	Val	Lys	Leu	Ala	Asp	Phe	Gly	Ser	Ala	Ser	Met	Ala	Ser	Pro	Ala	Asn	
					165					170					175		
	Ser	Phe	Val	Gly	Thr	Pro	Tyr	Trp	Met	Ala	Pro	Glu	Val	Ile	Leu	Ala	
35				180					185					190			
	Met	Asp	Glu	Gly	Gln	Tyr	Asp	Gly	Lys	Val	Asp	Ile	Trp	Ser	Leu	Gly	
		195						200					205				
40	Ile	Thr	Cys	Ile	Glu	Leu	Ala	Glu	Arg	Lys	Pro	Pro	Leu	Phe	Asn	Met	
		210					215					220					
	Asn	Ala	Met	Ser	Ala	Leu	Tyr	His	Ile	Ala	Gln	Asn	Asp	Ser	Pro	Thr	
	225					230					235					240	
45	Leu	Gln	Ser	Asn	Glu	Trp	Thr	Asp	Ser	Phe	Arg	Arg	Phe	Val	Asp	Tyr	
					245					250					255		
	Cys	Leu	Gln	Lys	Ile	Pro	Gln	Glu	Arg	Pro	Thr	Ser	Ala	Glu	Leu	Leu	
				260					265					270			
50	Arg	His	Asp	Phe	Val	Arg	Arg	Asp	Arg	Pro	Leu	Arg	Val	Leu	Ile	Asp	
			275					280					285				
55	Leu	Ile	Gln	Arg	Thr	Lys	Asp	Ala	Val	Arg	Glu	Leu	Asp	Asn	Leu	Gln	
		290					295					300					

	Tyr	Arg	Lys	Met	Lys	Lys	Ile	Leu	Phe	Gln	Glu	Thr	Arg	Asn	Gly	Pro	305	310	315	320
5	Leu	Asn	Glu	Ser	Gln	Glu	Asp	Glu	Glu	Asp	Ser	Glu	His	Gly	Thr	Ser	325	330	335	
	Leu	Asn	Arg	Glu	Met	Asp	Ser	Leu	Gly	Ser	Asn	His	Ser	Ile	Pro	Ser	340	345	350	
10	Met	Ser	Val	Ser	Thr	Gly	Ser	Gln	Ser	Ser	Ser	Val	Asn	Ser	Met	Gln	355	360	365	
	Glu	Val	Met	Asp	Glu	Ser	Ser	Ser	Glu	Leu	Val	Met	Met	His	Asp	Asp	370	375	380	
15	Glu	Ser	Thr	Ile	Asn	Ser	Ser	Ser	Ser	Val	Val	His	Lys	Lys	Asp	His	385	390	395	400
20	Val	Phe	Ile	Arg	Asp	Glu	Ala	Gly	His	Gly	Asp	Pro	Arg	Pro	Glu	Pro	405	410	415	
	Arg	Pro	Thr	Gln	Ser	Val	Gln	Ser	Gln	Ala	Leu	His	Tyr	Arg	Asn	Arg	420	425	430	
25	Glu	Arg	Phe	Ala	Thr	Ile	Lys	Ser	Ala	Ser	Leu	Val	Thr	Arg	Gln	Ile	435	440	445	
	His	Glu	His	Glu	Gln	Glu	Asn	Glu	Leu	Arg	Glu	Gln	Met	Ser	Gly	Tyr	450	455	460	
30	Lys	Arg	Met	Arg	Arg	Gln	His	Gln	Lys	Gln	Leu	Ile	Ala	Leu	Glu	Asn	465	470	475	480
	Lys	Leu	Lys	Ala	Glu	Met	Asp	Glu	His	Arg	Leu	Lys	Leu	Gln	Lys	Glu	485	490	495	
35	Val	Glu	Thr	His	Ala	Asn	Asn	Ser	Ser	Ile	Glu	Leu	Glu	Lys	Leu	Ala	500	505	510	
40	Lys	Lys	Gln	Val	Ala	Ile	Ile	Glu	Lys	Glu	Ala	Lys	Val	Ala	Ala	Ala	515	520	525	
	Asp	Glu	Lys	Lys	Phe	Gln	Gln	Gln	Ile	Leu	Ala	Gln	Gln	Lys	Lys	Asp	530	535	540	
45	Leu	Thr	Thr	Phe	Leu	Glu	Ser	Gln	Lys	Lys	Gln	Tyr	Lys	Ile	Cys	Lys	545	550	555	560
50	Glu	Lys	Ile	Lys	Glu	Glu	Met	Asn	Glu	Asp	His	Ser	Thr	Pro	Lys	Lys	565	570	575	
	Glu	Lys	Gln	Glu	Arg	Ile	Ser	Lys	His	Lys	Glu	Asn	Leu	Gln	His	Thr	580	585	590	
55	Gln	Ala	Glu	Glu	Glu	Ala	His	Leu	Leu	Thr	Gln	Gln	Arg	Leu	Tyr	Tyr	595	600	605	

[illegible]

<210> 5
<211> 4971
5 <212> DNA
<213> Homo sapiens
10 <300>
<308> Genbank NM_016151.1
15 <309> 2002-11-05
<313> (1) .. (4971)
20 <400> 5
aattcggcac gagctgagac ggagaagagg agaggcagag agggcgcggg gaccgtcagc 60
agcaccttag ctacaatcgt tcagctattc tcggaagaga gaagggagag ggaggaggcc 120
25 ggggcgggag tgggggctgt caccctcgga ccccggcgtg agaggggccc tgcgggccgga 180
cgctcctcggg gtgggcccc agtcggtggc cgaagacctc cagctcaggc ccctgggtcc 240
caaatttcca ggctttgccc ctctccttt ctcatagacc cgggtaacag tcctcatagt 300
30 ccagatatcc gggactcggg tcccaacctc tctaaacctg ggtctctgtt tcatagattt 360
tcaaatatca ggttcaggcc cctgcgtgca ccagtatccg gggttcatte cccgggctt 420
35 tcaaatatcg gattcagttc ccatcccggt cagatatcgg gggttcagac cccacaatca 480
gaaatccgga attcggcagc tgtcgccctc gacgaggggg aggactggac cgcgagggtca 540
gatttaggttg tcacccccctc ccctccaggg gaggtctccc gggcccgcgc ctcatggaag 600
40 gcgaaagccg aggaagaggt ggcaagggga aaggtctcct tgccccctctc cctgcttggc 660
agagccgctg gaggacccca ggcggaagcg gaggcgctgg ggcaccatag tgaccctac 720
45 caggccaggc cccactctca gggccccccag gggccaccat gccagctggg ggccgggccc 780
ggagcctgaa ggaccagat gtggctgagc tcttcttcaa ggatgaccca gaaaagctct 840
tctctgacct ccgggaaatt ggccatggca gctttggagc cgtatacttt gcccgggatg 900
50 tccggaatag tgaggtggtg gccatcaaga agatgtccta cagtgggaag cagtccaatg 960
agaaatggca agacatcatc aaggaggtgc ggttcttaca gaagctccgg catcccaaca 1020
55 ccattcagta ccggggctgt tacctgaggg agcacacggc ttggctggta atggagtatt 1080
gcctgggctc agcttctgac cttctagaag tgcacaagaa accccttcag gaggtagaga 1140

	tgcgagctgt	gacccacggg	gcgcttcagg	gcctggcata	tctgcactcc	cacaacatga	1200
5	tccatagggg	tgtgaaggct	ggaaacatcc	tgctgtcaga	gccaggggta	gtgaagctag	1260
	gggacttttg	ttctgcgtcc	atcatggcac	ctgccaaactc	cttcgtgggc	accccatact	1320
	ggatggcacc	cgaggtgatc	ctggccatgg	atgaggggca	gtacgatggc	aaagtggacg	1380
10	tctggtcctt	ggggataacc	tgcatcgagc	tggttgaacg	gaaaccaccg	ctctttaaca	1440
	tgaatgcat	gagtgcctta	taccacattg	cacagaacga	atccccctg	ctccagtcag	1500
15	gacactggtc	tgagtacttc	cggaattttg	tcgactcctg	tcttcagaaa	atccctcaag	1560
	acagaccaac	ctcagagggt	ctcctgaagc	accgctttgt	gctccgggag	cggccaccca	1620
	cagtcacat	ggacctgatc	cagaggacca	aggatgccgt	gcgggagctg	gacaacctgc	1680
20	agtaccgcaa	gatgaagaag	atcctgttcc	aagaggcacc	caacggccct	ggtgccgagg	1740
	ccccagagga	ggaagaggag	gccgagccct	acatgcaccg	ggccgggact	ctgaccagcc	1800
25	tcgagagtag	ccactcagtg	cccagcatgt	ccatcagcgc	ctccagccag	agcagctccg	1860
	tcaacagcct	agcagatgcc	tcagacaacg	aggaagagga	ggaggaggag	gaggaagagg	1920
	aggaggagga	agaaggccct	gaagcccggg	agatggccat	gatgcaggag	ggggagcaca	1980
30	cagtcacctc	tcacagctcc	attatccacc	ggctgccggg	ctctgacaac	ctatatgatg	2040
	accctacca	gccagagata	acccccagcc	ctctccagcc	gcctgcagcc	ccagctccca	2100
35	cttcaccac	ctcttcgcc	cgccgccggg	cctactgccg	taaccgagac	cactttgcc	2160
	ccatccgaac	cgctccctg	gtcagccgtc	agatccagga	gcatgagcag	gactctgcgc	2220
	tgccgggagca	gctgagcggc	tataagcggg	tgcgacgaca	gcaccagaag	cagctgctgg	2280
40	ccctggagtc	acggctgagg	ggtgaacggg	aggagcacag	tgcacggctg	cagcgggagc	2340
	ttgaggcgca	gcgggctggc	tttggggcag	aggcagaaaa	gctggcccgg	cggcaccagg	2400
45	ccataggtga	gaaggaggca	cgagctgcc	aggccgagga	gcggaagtgc	cagcagcaca	2460
	tccttgggca	gcagaagaag	gagctggctg	ccctgctgga	ggcacagaag	cggacctaca	2520
	aacttcgcaa	ggaacagctg	aaggaggagc	tccaggagaa	ccccagcact	ccaagcggg	2580
50	agaaggccga	gtggctgctg	cggcagaagg	agcagctcca	gcagtgccag	gcggaggagg	2640
	aagcagggct	gctgcggcgg	cagcgccagt	actttgagct	gcagtgtcgc	cagtacaagc	2700
55	gcaagatgtt	gctggctcgg	cacagcctgg	accaggacct	gctgcgggag	gacctgaaca	2760
	agaagcagac	ccagaaggac	ttggagtgtg	cactgctgct	tcggcagcac	gaggccacgc	2820

gggagctgga gctgcggcag ctccaggccg tgcagcgcac gcgggctgag ctcacccgcc 2880
tgcagcacca gacggagctg ggcaaccagc tggagtacaa caagcggcgt gagcaagagt 2940
5 tgcggcagaa gcatgcggcc cagggttcgcc agcagcccaa gagcctcaaa gtacgtgcag 3000
gccagcgcgc cccgggcctt ccactcccca ttcttggggc tctggggcca cccaacacag 3060
gcacccctat agaacagcag ccctgctcac ctggccagga ggagtcctg gaccaaagaa 3120
10 tgcttggcga ggaggaggaa gcagttggag agagaaggat tctgggaaag gaagggggcca 3180
ctttggagcc caagcagcag aggattctgg gggagaatac aggagcccct agtcccagtc 3240
15 cacaaaaaca tgggagcctg gttgatgagg aagtttgggg tctgcctgag gagatagagg 3300
agcttagggg gccctccctt gtaccccagg agaggagcat tgttggccag gaggaggctg 3360
ggacgtggag cttgtggggg aaggaggatg agagtcttct ggatgaggag tttgagcttg 3420
20 gctgggtcca gggcccagca ctgactcccg tccctgagga ggaggaagaa gaggaagagg 3480
gggctccgat tgggacccct agggatcctg gagatggttg tccttcccc cactccctc 3540
25 ctgaaccccc tccaacacac ctgaggccct gccctgccag ccagctccct ggactcctgt 3600
cccatggcct cctggccggc ctctcctttg cagtggggtc ctctctggc ctctgcccc 3660
tctgtctgct gctgtctgtt ccattgctgg cagcccaggg tgggggtggc ctgcaggcag 3720
30 cgctgctggc ccttgagggtg gggctggtgg gtctgggggc ctctacctg ctctttgtg 3780
cagccctgca cctgcccctc agtcttttcc tactcctggc ccagggtacc gactggggg 3840
35 ccgtcctggg cctgagctgg cgccgaggcc tcatgggtgt tcccctgggc cttggagctg 3900
cctggctctt agcttgggca ggcctagctc tacctctggt ggctatggca gcggggggca 3960
gatgggtgcg gcagcagggc cccgggtgct gccggggcat atctcgactc tggttgcggg 4020
40 ttctgctgcg cctgtcacc ctaggccttcc gggccctgca gggctgtggg gctgtggggg 4080
accggggtct gtttgactg taccctaaaa ccaacaagga tggcttccgc agccgcctgc 4140
45 ccgtccctgg gcccggcg cgtaatcccc gcaccacca acaccatta gctctgttgg 4200
caagggtctg ggtcctgtgc aagggtgga actggcgtct ggcacgggac agccagggtt 4260
tagcatccca cttgcccccg tgggccatcc acacactggc cagctggggc ctgcttcggg 4320
50 gtgaacggcc caccgaatc cccgggtac taccacgag ccagcgccag ctaggggccc 4380
ctgcctccca ccagccactg ccagggactc tagccggcg gaggtcacgc acccgccagt 4440
55 cccgggcctt gccccctgg aggtagctga ctccagccct tccagcccaa atctagagca 4500
ttgagcactt tatctccac gactcagtga agtttctcca gtccctagtc ctctcttttc 4560


```

accacaccttc ctcagtttgc tcacttaccc caggcccagc ccttcggacc tctagacagg 4620
cagcctcctc agctgtggag tccagcagtc actctgtgtt ctctgggcgc tcctccccta 4680
5 agttattgct gttcgccccgc tgtgtgtgct catcctcacc ctcattgact caggcctggg 4740
gccagggggtg gtggaggggtg ggaagagtca tgtttttttt ctctcttttg attttgtttt 4800
10 tctgtctccc ttccaacctg tccccttccc cccaccaaaa aaagaaaaag acaaacacaa 4860
ataaaatata tgagcggaac tgtgaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 4920
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 4971
15

```

```

<210> 6
<211> 1235
20 <212> PRT
<213> Homo sapiens

```

```

25 <300>
<308> Genbank NP_057235.1
30 <309> 2002-11-05
<313> (1)..(1235)

```

```

35 <400> 6
Met Pro Ala Gly Gly Arg Ala Gly Ser Leu Lys Asp Pro Asp Val Ala
1 5 10 15
40 Glu Leu Phe Phe Lys Asp Asp Pro Glu Lys Leu Phe Ser Asp Leu Arg
20 25 30
Glu Ile Gly His Gly Ser Phe Gly Ala Val Tyr Phe Ala Arg Asp Val
35 40 45
45 Arg Asn Ser Glu Val Val Ala Ile Lys Lys Met Ser Tyr Ser Gly Lys
50 55 60
Gln Ser Asn Glu Lys Trp Gln Asp Ile Ile Lys Glu Val Arg Phe Leu
50 65 70 75 80
Gln Lys Leu Arg His Pro Asn Thr Ile Gln Tyr Arg Gly Cys Tyr Leu
85 90 95
55 Arg Glu His Thr Ala Trp Leu Val Met Glu Tyr Cys Leu Gly Ser Ala
100 105 110

```

	Ser	Asp	Leu	Leu	Glu	Val	His	Lys	Lys	Pro	Leu	Gln	Glu	Val	Glu	Ile	
			115					120					125				
5	Ala	Ala	Val	Thr	His	Gly	Ala	Leu	Gln	Gly	Leu	Ala	Tyr	Leu	His	Ser	
			130				135					140					
	His	Asn	Met	Ile	His	Arg	Asp	Val	Lys	Ala	Gly	Asn	Ile	Leu	Leu	Ser	
	145					150					155					160	
10	Glu	Pro	Gly	Leu	Val	Lys	Leu	Gly	Asp	Phe	Gly	Ser	Ala	Ser	Ile	Met	
				165						170						175	
	Ala	Pro	Ala	Asn	Ser	Phe	Val	Gly	Thr	Pro	Tyr	Trp	Met	Ala	Pro	Glu	
15				180					185					190			
	Val	Ile	Leu	Ala	Met	Asp	Glu	Gly	Gln	Tyr	Asp	Gly	Lys	Val	Asp	Val	
			195					200					205				
20	Trp	Ser	Leu	Gly	Ile	Thr	Cys	Ile	Glu	Leu	Ala	Glu	Arg	Lys	Pro	Pro	
	210						215					220					
	Leu	Phe	Asn	Met	Asn	Ala	Met	Ser	Ala	Leu	Tyr	His	Ile	Ala	Gln	Asn	
	225					230					235					240	
25	Glu	Ser	Pro	Val	Leu	Gln	Ser	Gly	His	Trp	Ser	Glu	Tyr	Phe	Arg	Asn	
				245						250					255		
	Phe	Val	Asp	Ser	Cys	Leu	Gln	Lys	Ile	Pro	Gln	Asp	Arg	Pro	Thr	Ser	
30				260					265					270			
	Glu	Val	Leu	Leu	Lys	His	Arg	Phe	Val	Leu	Arg	Glu	Arg	Pro	Pro	Thr	
			275					280					285				
35	Val	Ile	Met	Asp	Leu	Ile	Gln	Arg	Thr	Lys	Asp	Ala	Val	Arg	Glu	Leu	
		290					295					300					
	Asp	Asn	Leu	Gln	Tyr	Arg	Lys	Met	Lys	Lys	Ile	Leu	Phe	Gln	Glu	Ala	
	305					310					315					320	
40	Pro	Asn	Gly	Pro	Gly	Ala	Glu	Ala	Pro	Glu	Glu	Glu	Glu	Glu	Ala	Glu	
					325					330					335		
	Pro	Tyr	Met	His	Arg	Ala	Gly	Thr	Leu	Thr	Ser	Leu	Glu	Ser	Ser	His	
45				340					345					350			
	Ser	Val	Pro	Ser	Met	Ser	Ile	Ser	Ala	Ser	Ser	Gln	Ser	Ser	Ser	Val	
			355					360					365				
50	Asn	Ser	Leu	Ala	Asp	Ala	Ser	Asp	Asn	Glu	Glu	Glu	Glu	Glu	Glu	Glu	
		370					375					380					
	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Gly	Pro	Glu	Ala	Arg	Glu	Met	Ala		
					390					395					400		
55	Met	Met	Gln	Glu	Gly	Glu	His	Thr	Val	Thr	Ser	His	Ser	Ser	Ile	Ile	
					405					410					415		

His Arg Leu Pro Gly Ser Asp Asn Leu Tyr Asp Asp Pro Tyr Gln Pro
 420 425 430
 5 Glu Ile Thr Pro Ser Pro Leu Gln Pro Pro Ala Ala Pro Ala Pro Thr
 435 440 445
 Ser Thr Thr Ser Ser Ala Arg Arg Arg Ala Tyr Cys Arg Asn Arg Asp
 450 455 460
 10 His Phe Ala Thr Ile Arg Thr Ala Ser Leu Val Ser Arg Gln Ile Gln
 465 470 475 480
 Glu His Glu Gln Asp Ser Ala Leu Arg Glu Gln Leu Ser Gly Tyr Lys
 485 490 495
 15 Arg Met Arg Arg Gln His Gln Lys Gln Leu Leu Ala Leu Glu Ser Arg
 500 505 510
 20 Leu Arg Gly Glu Arg Glu Glu His Ser Ala Arg Leu Gln Arg Glu Leu
 515 520 525
 Glu Ala Gln Arg Ala Gly Phe Gly Ala Glu Ala Glu Lys Leu Ala Arg
 530 535 540
 25 Arg His Gln Ala Ile Gly Glu Lys Glu Ala Arg Ala Ala Gln Ala Glu
 545 550 555 560
 Glu Arg Lys Phe Gln Gln His Ile Leu Gly Gln Gln Lys Lys Glu Leu
 565 570 575
 30 Ala Ala Leu Leu Glu Ala Gln Lys Arg Thr Tyr Lys Leu Arg Lys Glu
 580 585 590
 35 Gln Leu Lys Glu Glu Leu Gln Glu Asn Pro Ser Thr Pro Lys Arg Glu
 595 600 605
 Lys Ala Glu Trp Leu Leu Arg Gln Lys Glu Gln Leu Gln Gln Cys Gln
 610 615 620
 40 Ala Glu Glu Glu Ala Gly Leu Leu Arg Arg Gln Arg Gln Tyr Phe Glu
 625 630 635 640
 Leu Gln Cys Arg Gln Tyr Lys Arg Lys Met Leu Leu Ala Arg His Ser
 645 650 655
 45 Leu Asp Gln Asp Leu Leu Arg Glu Asp Leu Asn Lys Lys Gln Thr Gln
 660 665 670
 50 Lys Asp Leu Glu Cys Ala Leu Leu Leu Arg Gln His Glu Ala Thr Arg
 675 680 685
 Glu Leu Glu Leu Arg Gln Leu Gln Ala Val Gln Arg Thr Arg Ala Glu
 690 695 700
 55 Leu Thr Arg Leu Gln His Gln Thr Glu Leu Gly Asn Gln Leu Glu Tyr
 705 710 715 720

	Asn	Lys	Arg	Arg	Glu	Gln	Glu	Leu	Arg	Gln	Lys	His	Ala	Ala	Gln	Val	
					725					730					735		
5	Arg	Gln	Gln	Pro	Lys	Ser	Leu	Lys	Val	Arg	Ala	Gly	Gln	Arg	Pro	Pro	
				740					745					750			
	Gly	Leu	Pro	Leu	Pro	Ile	Pro	Gly	Ala	Leu	Gly	Pro	Pro	Asn	Thr	Gly	
			755					760					765				
10	Thr	Pro	Ile	Glu	Gln	Gln	Pro	Cys	Ser	Pro	Gly	Gln	Glu	Ala	Val	Leu	
		770					775					780					
	Asp	Gln	Arg	Met	Leu	Gly	Glu	Glu	Glu	Glu	Ala	Val	Gly	Glu	Arg	Arg	
	785					790					795					800	
15	Ile	Leu	Gly	Lys	Glu	Gly	Ala	Thr	Leu	Glu	Pro	Lys	Gln	Gln	Arg	Ile	
					805					810					815		
	Leu	Gly	Glu	Glu	Ser	Gly	Ala	Pro	Ser	Pro	Ser	Pro	Gln	Lys	His	Gly	
20				820					825					830			
	Ser	Leu	Val	Asp	Glu	Glu	Val	Trp	Gly	Leu	Pro	Glu	Glu	Ile	Glu	Glu	
			835					840					845				
25	Leu	Arg	Val	Pro	Ser	Leu	Val	Pro	Gln	Glu	Arg	Ser	Ile	Val	Gly	Gln	
		850					855					860					
	Glu	Glu	Ala	Gly	Thr	Trp	Ser	Leu	Trp	Gly	Lys	Glu	Asp	Glu	Ser	Leu	
	865					870					875					880	
30	Leu	Asp	Glu	Glu	Phe	Glu	Leu	Gly	Trp	Val	Gln	Gly	Pro	Ala	Leu	Thr	
					885					890					895		
	Pro	Val	Pro	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Gly	Ala	Pro	Ile	Gly	
35				900					905					910			
	Thr	Pro	Arg	Asp	Pro	Gly	Asp	Gly	Cys	Pro	Ser	Pro	Asp	Ile	Pro	Pro	
			915					920					925				
40	Glu	Pro	Pro	Pro	Thr	His	Leu	Arg	Pro	Cys	Pro	Ala	Ser	Gln	Leu	Pro	
		930					935					940					
	Gly	Leu	Leu	Ser	His	Gly	Leu	Leu	Ala	Gly	Leu	Ser	Phe	Ala	Val	Gly	
	945					950					955					960	
45	Ser	Ser	Ser	Gly	Leu	Leu	Pro	Leu	Leu	Leu	Leu	Leu	Leu	Leu	Pro	Leu	
					965				970						975		
	Leu	Ala	Ala	Gln	Gly	Gly	Gly	Gly	Leu	Gln	Ala	Ala	Leu	Leu	Ala	Leu	
50				980					985					990			
	Glu	Val	Gly	Leu	Val	Gly	Leu	Gly	Ala	Ser	Tyr	Leu	Leu	Leu	Cys	Thr	
			995				1000						1005				
55	Ala	Leu	His	Leu	Pro	Ser	Ser	Leu	Phe	Leu	Leu	Leu	Ala	Gln	Gly		
		1010					1015						1020				

Thr Ala Leu Gly Ala Val Leu Gly Leu Ser Trp Arg Arg Gly Leu
 1025 1030 1035
 5 Met Gly Val Pro Leu Gly Leu Gly Ala Ala Trp Leu Leu Ala Trp
 1040 1045 1050
 Pro Gly Leu Ala Leu Pro Leu Val Ala Met Ala Ala Gly Gly Arg
 1055 1060 1065
 10 Trp Val Arg Gln Gln Gly Pro Arg Val Arg Arg Gly Ile Ser Arg
 1070 1075 1080
 Leu Trp Leu Arg Val Leu Leu Arg Leu Ser Pro Met Ala Phe Arg
 1085 1090 1095
 15 Ala Leu Gln Gly Cys Gly Ala Val Gly Asp Arg Gly Leu Phe Ala
 1100 1105 1110
 Leu Tyr Pro Lys Thr Asn Lys Asp Gly Phe Arg Ser Arg Leu Pro
 1115 1120 1125
 Val Pro Gly Pro Arg Arg Arg Asn Pro Arg Thr Thr Gln His Pro
 1130 1135 1140
 25 Leu Ala Leu Leu Ala Arg Val Trp Val Leu Cys Lys Gly Trp Asn
 1145 1150 1155
 Trp Arg Leu Ala Arg Ala Ser Gln Gly Leu Ala Ser His Leu Pro
 1160 1165 1170
 30 Pro Trp Ala Ile His Thr Leu Ala Ser Trp Gly Leu Leu Arg Gly
 1175 1180 1185
 Glu Arg Pro Thr Arg Ile Pro Arg Leu Leu Pro Arg Ser Gln Arg
 1190 1195 1200
 Gln Leu Gly Pro Pro Ala Ser His Gln Pro Leu Pro Gly Thr Leu
 1205 1210 1215
 40 Ala Gly Arg Arg Ser Arg Thr Arg Gln Ser Arg Ala Leu Pro Pro
 1220 1225 1230
 Trp Arg
 1235
 45
 <210> 7
 <211> 4242
 50
 <212> DNA
 <213> Homo sapiens
 55
 <300>

<308> Genbank NM_004783

<309> 2000-11-01

5 <313> (1) .. (4242)

<400> 7

10	agaatttcaa atatcagggt caggcccctg cgtgcaccag tatccggggt tcattccccg	60
	ggcgttcaaa tatcggattc agtctccatc ccgttcagat attcgggggt cagacccac	120
	aatcagaaat ccggaattcg gcagctgtcg ccctcgacga gggggaggac tggaccgca	180
15	ggtcagatta ggttgtcacc ccctcccctc caggggaggc ttcccgggcc cgcccctcag	240
	gaagggcgaa agccgaggaa gaggtggcaa ggggaaaggt ctcttgccc ctctccctgc	300
20	ttggcagagc cgctggagga cccagggcgg aagcggaggc gctggggcac catagtgacc	360
	cctaccaggc caggccccac tctcagggcc ccaggggcc accatgccag ctggggggccg	420
	ggccgggagc ctgaaggacc cagatgtggc tgagctcttc ttcaaggatg acccagaaaa	480
25	gctcttctct gacctccggg aaattggcca tggcagcttt ggagccgtat actttgccc	540
	ggatgtccgg aatagtgagg tgggtggcat caagaagatg tcctacagtg ggaagcagtc	600
30	caatgagaaa tggcaagaca tcatcaagga ggtgcggttc ttacagaagc tccggcatcc	660
	caacaccatt cagtaccggg gctgttacct gagggagcac acggcttggc tggtaatgga	720
	gtattgcctg ggctcagctt ctgaccttct agaagtgcac aagaaacccc ttcaggaggt	780
35	agagatcgca gctgtgacct acggggcgct tcagggcctg gcatatctgc actcccacaa	840
	catgatccat agggatgtga aggttggaat catcctgctg tcagagccag ggttagtgaa	900
40	gctaggggac tttggttctg cgtccatcat ggcacctgcc aactccttcg tgggcacccc	960
	atactggatg gcacccgagg tgatcctggc catggatgag gggcagtagc atggcaaagt	1020
	ggacgtctgg tccttgggga taacctgcat cgagctggct gaacggaaac caccgctctt	1080
45	taacatgaat gcgatgagt ccttatacca cattgcacag aacgaatccc ccgtgctcca	1140
	gtcaggacac tgggtctgagt acttccggaa ttttgtcgac tcctgtcttc agaaaatccc	1200
50	tcaagacaga ccaacctcag aggttctcct gaagcaccgc tttgtgctcc gggagcggcc	1260
	accacagtc atcatggacc tgatccagag gaccaaggat gccgtgcggg agctggacaa	1320
	cctgcagtac cgcaagatga agaagatcct gttccaagag gcacccaacg gccctggtgc	1380
55	cgaggcccca gaggaggaag aggagggcga gccctacatg caccggggcc ggactctgac	1440
	cagcctcgag agtagccact cagtggccag catgtccatc agcgcctcca gccagagcag	1500

ctccgtcaac agcctagcag atgcctcaga caacgaggaa gaggaggagg aggaggagga 1560
5 agaggaggag gaggaagaag gccctgaagc ccgggagatg gccatgatgc aggaggggga 1620
gcacacagtc acctctcaca gctccattat ccaccggctg ccgggctctg acaacctata 1680
tgatgacccc taccagccag agataacccc cagccctctc cagccgcctg cagccccagc 1740
10 tcccacttcc accacctctt ctgcccgcg ccgggcctac tgccgtaacc gagaccactt 1800
tgccaccatc cgaaccgcct ccctggtcag ccgtcagatc caggagcatg agcaggactc 1860
tgcgctgcgg gagcagctga gcggtataa gcggatgcga cgacagcacc agaagcagct 1920
15 gctggccctg gagtcacggc tgaggggtga acgggaggag cacagtgcac ggctgcagcg 1980
ggagcttgag gcgcagcggg ctggctttgg ggcagaggca gaaaagctgg cccggcggca 2040
20 ccaggccata ggtgagaagg aggcacgagc tgcccaggcc gaggagcggg agttccagca 2100
gcacatectt gggcagcaga agaaggagct ggctgcctg ctggaggcac agaagcggac 2160
ctacaaactt cgcaaggaa agctgaagga ggagctccag gagaacccca gcactcccaa 2220
25 gcgggagaag gccgagtggc tgctgcggca gaaggagcag ctccagcagt gccaggcggg 2280
ggaggaagca gggctgctgc ggcggcagcg ccagtacttt gagctgcagt gtcgccagta 2340
30 caagcgcaag atgttgctgg ctcggcacag cctggaccag gacctgctgc gggaggacct 2400
gaacaagaag cagaccaga aggacttggg gtgtgcactg ctgcttcggc agcacgaggc 2460
cacgcgggag ctggagctgc ggcagctcca ggccgtgcag cgcacgcggg ctgagctcac 2520
35 ccgcctgcag caccagacgg agctgggcaa ccagctggag tacaacaagc ggcgtgagca 2580
agagttgcgg cagaagcatg cggcccaggt tcgccagcag cccaagagcc tcaaactaa 2640
40 ggagctgcag atcaagaagc agttccagga gacgtgtaag atccagactc ggcagtacaa 2700
ggctctgcga gcacacttgc tggagaccac gcccaaagct cagcacaaga gcctccttaa 2760
gcggctcaag gaagagcaga cccgcaagct ggcgatcttg gcggagcagt atgaccagtc 2820
45 catctcagag atgctcagct cacaggcgct gcggcttgat gagaccagg aggcagagtt 2880
ccaggccctt cggcagcagc ttcaacagga gctggagctg ctcaacgctt accagagcaa 2940
50 gatcaagatc cgcacagaga gccagacga gaggagctg cgggagctgg agcagagggt 3000
cgcgctgcgg cgggcactgc tggagcagcg ggtggaagag gagctgctgg ccctgcagac 3060
aggacgctcc gagcgaatcc gcagtctgct tgagcggcag gcccgtaga tcgaggcctt 3120
55 cgatgcggaa agcatgaggc tgggcttctc cagcatggct ctggggggca tcccggctga 3180

agctgctgcc cagggctatc ctgctccacc ccctgcccc aacctggccct ccggtcccg 3240
tccccgttct ggggcacact ggagccatgg ccctcctcca ccaggcatgc cccctccagc 3300
5 ctggcgctcag ccgtctctgc tggctcccc agggcccca aactggctgg gggcccccac 3360
acaaagtggg acaccccggt ggggagccct gctgctgcta agaaacagcc cccagccctt 3420
10 ggggggggca gcctcggggg gcagtggcag tgagaatgtg gggcccccctg ctgccgcggt 3480
ggccggggccc ctgagccgca gcaccagtgt cgcttcccc atcctcaatg gttcttccca 3540
cttctattcc tgaggtgcag cggggaggag cagatgagct gggcagggca ggggtgggtg 3600
15 gagcctgacc ctggagggca ctgagctgga gggccctgca agggtagggg acaagatgta 3660
ggctccagct cccctcagac ctctcatct catgagcttc ttggggctgg ccagtggccc 3720
agggccagct tggcgataga tgctcaagg ctgcctggga gggcgctc cctaccatgg 3780
20 tgccaggggt ctccctccgc cacctaggaa aggagggaga tgtgctgtc aaatattcat 3840
ctagtcctct gggggagggg aagggtgggt ctagacatac tatattcaga gaactatact 3900
25 accctcacag tgaggccctc agacctgcca cagggcagag caggtctggg gcctgaggca 3960
gggagaatga gaggccacct tactggcagg aaggatcagg atggggtctt ggggtcagga 4020
tgcttgggtc tcttccgta actgtctgac gtcctgtgcc gtcttgcct ttatcttttt 4080
30 tttttttttt taattgggat cagggtctgg ggggggaaac aagggaagga ccttggaagg 4140
ggctgctccc aggcctgggg ggcagtcgtg ggagccctc tcagctgtgg ggctggcaca 4200
35 gagccccagg caagctttta ataaactgtt gggtattcta ac 4242

<210> 8

40 <211> 1049

<212> PRT

45 <213> Homo sapiens

<300>

50 <308> Genbank NP_004774.1

<309> 2000-11-01

<313> (1)..(1049)

55

<400> 8

Met Pro Ala Gly Gly Arg Ala Gly Ser Leu Lys Asp Pro Asp Val Ala
 1 5 10 15
 5 Glu Leu Phe Phe Lys Asp Asp Pro Glu Lys Leu Phe Ser Asp Leu Arg
 20 25 30
 Glu Ile Gly His Gly Ser Phe Gly Ala Val Tyr Phe Ala Arg Asp Val
 35 40 45
 10 Arg Asn Ser Glu Val Val Ala Ile Lys Lys Met Ser Tyr Ser Gly Lys
 50 55 60
 Gln Ser Asn Glu Lys Trp Gln Asp Ile Ile Lys Glu Val Arg Phe Leu
 65 70 75 80
 15 Gln Lys Leu Arg His Pro Asn Thr Ile Gln Tyr Arg Gly Cys Tyr Leu
 85 90 95
 Arg Glu His Thr Ala Trp Leu Val Met Glu Tyr Cys Leu Gly Ser Ala
 100 105 110
 Ser Asp Leu Leu Glu Val His Lys Lys Pro Leu Gln Glu Val Glu Ile
 115 120 125
 25 Ala Ala Val Thr His Gly Ala Leu Gln Gly Leu Ala Tyr Leu His Ser
 130 135 140
 His Asn Met Ile His Arg Asp Val Lys Ala Gly Asn Ile Leu Leu Ser
 145 150 155 160
 30 Glu Pro Gly Leu Val Lys Leu Gly Asp Phe Gly Ser Ala Ser Ile Met
 165 170 175
 Ala Pro Ala Asn Ser Phe Val Gly Thr Pro Tyr Trp Met Ala Pro Glu
 180 185 190
 Val Ile Leu Ala Met Asp Glu Gly Gln Tyr Asp Gly Lys Val Asp Val
 195 200 205
 40 Trp Ser Leu Gly Ile Thr Cys Ile Glu Leu Ala Glu Arg Lys Pro Pro
 210 215 220
 Leu Phe Asn Met Asn Ala Met Ser Ala Leu Tyr His Ile Ala Gln Asn
 225 230 235 240
 45 Glu Ser Pro Val Leu Gln Ser Gly His Trp Ser Glu Tyr Phe Arg Asn
 245 250 255
 Phe Val Asp Ser Cys Leu Gln Lys Ile Pro Gln Asp Arg Pro Thr Ser
 260 265 270
 Glu Val Leu Leu Lys His Arg Phe Val Leu Arg Glu Arg Pro Pro Thr
 275 280 285
 55 Val Ile Met Asp Leu Ile Gln Arg Thr Lys Asp Ala Val Arg Glu Leu
 290 295 300

	Asp	Asn	Leu	Gln	Tyr	Arg	Lys	Met	Lys	Lys	Ile	Leu	Phe	Gln	Glu	Ala	305	310	315	320
5	Pro	Asn	Gly	Pro	Gly	Ala	Glu	Ala	Pro	Glu	Glu	Glu	Glu	Glu	Ala	Glu	325	330	335	
	Pro	Tyr	Met	His	Arg	Ala	Gly	Thr	Leu	Thr	Ser	Leu	Glu	Ser	Ser	His	340	345	350	
10	Ser	Val	Pro	Ser	Met	Ser	Ile	Ser	Ala	Ser	Ser	Gln	Ser	Ser	Ser	Val	355	360	365	
	Asn	Ser	Leu	Ala	Asp	Ala	Ser	Asp	Asn	Glu	Glu	Glu	Glu	Glu	Glu	Glu	370	375	380	
15	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Gly	Pro	Glu	Ala	Arg	Glu	Met	Ala	385	390	395	400
	Met	Met	Gln	Glu	Gly	Glu	His	Thr	Val	Thr	Ser	His	Ser	Ser	Ile	Ile	405	410	415	
20	His	Arg	Leu	Pro	Gly	Ser	Asp	Asn	Leu	Tyr	Asp	Asp	Pro	Tyr	Gln	Pro	420	425	430	
	Glu	Ile	Thr	Pro	Ser	Pro	Leu	Gln	Pro	Pro	Ala	Ala	Pro	Ala	Pro	Thr	435	440	445	
25	Ser	Thr	Thr	Ser	Ser	Ala	Arg	Arg	Arg	Ala	Tyr	Cys	Arg	Asn	Arg	Asp	450	455	460	
30	His	Phe	Ala	Thr	Ile	Arg	Thr	Ala	Ser	Leu	Val	Ser	Arg	Gln	Ile	Gln	465	470	475	480
	Glu	His	Glu	Gln	Asp	Ser	Ala	Leu	Arg	Glu	Gln	Leu	Ser	Gly	Tyr	Lys	485	490	495	
35	Arg	Met	Arg	Arg	Gln	His	Gln	Lys	Gln	Leu	Leu	Ala	Leu	Glu	Ser	Arg	500	505	510	
40	Leu	Arg	Gly	Glu	Arg	Glu	Glu	His	Ser	Ala	Arg	Leu	Gln	Arg	Glu	Leu	515	520	525	
	Glu	Ala	Gln	Arg	Ala	Gly	Phe	Gly	Ala	Glu	Ala	Glu	Lys	Leu	Ala	Arg	530	535	540	
45	Arg	His	Gln	Ala	Ile	Gly	Glu	Lys	Glu	Ala	Arg	Ala	Ala	Gln	Ala	Glu	545	550	555	560
	Glu	Arg	Lys	Phe	Gln	Gln	His	Ile	Leu	Gly	Gln	Gln	Lys	Lys	Glu	Leu	565	570	575	
50	Ala	Ala	Leu	Leu	Glu	Ala	Gln	Lys	Arg	Thr	Tyr	Lys	Leu	Arg	Lys	Glu	580	585	590	
55	Gln	Leu	Lys	Glu	Glu	Leu	Gln	Glu	Asn	Pro	Ser	Thr	Pro	Lys	Arg	Glu	595	600	605	

	Lys	Ala	Glu	Trp	Leu	Leu	Arg	Gln	Lys	Glu	Gln	Leu	Gln	Gln	Cys	Gln	
	610						615					620					
5	Ala	Glu	Glu	Glu	Ala	Gly	Leu	Leu	Arg	Arg	Gln	Arg	Gln	Tyr	Phe	Glu	
	625					630					635					640	
	Leu	Gln	Cys	Arg	Gln	Tyr	Lys	Arg	Lys	Met	Leu	Leu	Ala	Arg	His	Ser	
					645					650					655		
10	Leu	Asp	Gln	Asp	Leu	Leu	Arg	Glu	Asp	Leu	Asn	Lys	Lys	Gln	Thr	Gln	
				660					665						670		
	Lys	Asp	Leu	Glu	Cys	Ala	Leu	Leu	Leu	Arg	Gln	His	Glu	Ala	Thr	Arg	
				675				680						685			
15	Glu	Leu	Glu	Leu	Arg	Gln	Leu	Gln	Ala	Val	Gln	Arg	Thr	Arg	Ala	Glu	
		690					695					700					
	Leu	Thr	Arg	Leu	Gln	His	Gln	Thr	Glu	Leu	Gly	Asn	Gln	Leu	Glu	Tyr	
20		705				710					715					720	
	Asn	Lys	Arg	Arg	Glu	Gln	Glu	Leu	Arg	Gln	Lys	His	Ala	Ala	Gln	Val	
					725					730					735		
25	Arg	Gln	Gln	Pro	Lys	Ser	Leu	Lys	Ser	Lys	Glu	Leu	Gln	Ile	Lys	Lys	
				740					745						750		
	Gln	Phe	Gln	Glu	Thr	Cys	Lys	Ile	Gln	Thr	Arg	Gln	Tyr	Lys	Ala	Leu	
			755					760					765				
30	Arg	Ala	His	Leu	Leu	Glu	Thr	Thr	Pro	Lys	Ala	Gln	His	Lys	Ser	Leu	
		770					775					780					
	Leu	Lys	Arg	Leu	Lys	Glu	Glu	Gln	Thr	Arg	Lys	Leu	Ala	Ile	Leu	Ala	
35		785				790					795					800	
	Glu	Gln	Tyr	Asp	Gln	Ser	Ile	Ser	Glu	Met	Leu	Ser	Ser	Gln	Ala	Leu	
					805					810					815		
40	Arg	Leu	Asp	Glu	Thr	Gln	Glu	Ala	Glu	Phe	Gln	Ala	Leu	Arg	Gln	Gln	
				820					825					830			
	Leu	Gln	Gln	Glu	Leu	Glu	Leu	Leu	Asn	Ala	Tyr	Gln	Ser	Lys	Ile	Lys	
			835					840					845				
45	Ile	Arg	Thr	Glu	Ser	Gln	His	Glu	Arg	Glu	Leu	Arg	Glu	Leu	Glu	Gln	
		850					855					860					
	Arg	Val	Ala	Leu	Arg	Arg	Ala	Leu	Leu	Glu	Gln	Arg	Val	Glu	Glu	Glu	
50		865				870					875					880	
	Leu	Leu	Ala	Leu	Gln	Thr	Gly	Arg	Ser	Glu	Arg	Ile	Arg	Ser	Leu	Leu	
					885					890					895		
55	Glu	Arg	Gln	Ala	Arg	Glu	Ile	Glu	Ala	Phe	Asp	Ala	Glu	Ser	Met	Arg	
				900					905						910		

Leu Gly Phe Ser Ser Met Ala Leu Gly Gly Ile Pro Ala Glu Ala Ala
915 920 925

5 Ala Gln Gly Tyr Pro Ala Pro Pro Pro Ala Pro Ala Trp Pro Ser Arg
930 935 940

Pro Val Pro Arg Ser Gly Ala His Trp Ser His Gly Pro Pro Pro Pro
945 950 955 960

10 Gly Met Pro Pro Pro Ala Trp Arg Gln Pro Ser Leu Leu Ala Pro Pro
965 970 975

Gly Pro Pro Asn Trp Leu Gly Pro Pro Thr Gln Ser Gly Thr Pro Arg
980 985 990

15 Gly Gly Ala Leu Leu Leu Leu Arg Asn Ser Pro Gln Pro Leu Arg Arg
995 1000 1005

Ala Ala Ser Gly Gly Ser Gly Ser Glu Asn Val Gly Pro Pro Ala
1010 1015 1020

Ala Ala Val Pro Gly Pro Leu Ser Arg Ser Thr Ser Val Ala Ser
1025 1030 1035

25 His Ile Leu Asn Gly Ser Ser His Phe Tyr Ser
1040 1045

30 <210> 9

<211> 3285

<212> DNA

35 <213> Homo sapiens

<220>

40 <221> misc_feature

<222> (3245)..(3245)

<223> a, t, c or g

45 <220>

<221> misc_feature

50 <222> (3263)..(3263)

<223> a, t, c or g

55 <220>

```

<221> misc_feature
<222> (3278)..(3278)
5 <223> a, t, c or g

<220>
10 <221> misc_feature
<222> (72)..(72)
<223> a, t, c or g
15
<300>
<308> EMBL AY049015
20 <309> 2001-10-15
<313> (1)..(3285)
25
<400> 9
caacggattt catttcatac agatgaacca aggatcgga tagcagtata aaattagaat 60
caagacagct gntcgccaag caggatgcca tcaactaaca gagcaggcag cctgaaggac 120
30 cctgaaattg cagagctctt cttcaaagaa gatccagaga agctcttcac agatctcaga 180
gaaattggcc atggaagctt tggagcagtg tattttgcac gagatgtgcg taccaatgaa 240
35 gtggtggcca tcaagaaaat gtcttatagt ggaaagcagt ctactgagaa atggcaggat 300
attattaagg aagtcaagtt tctacaaaga ataaaacatc ccaacagtat agaatacaaa 360
ggctgttatt tacgtgaaca cacagcatgg cttgtaatgg aatattgttt aggatctgct 420
40 tcggatttac tagaagttca caaaaagcca ttacaagaag tggaaatagc agcaattaca 480
catggtgctc ttcagggatt agcctactta cattctcata ctatgattca tagagatatc 540
45 aaagcaggaa atatccttct gacagaacca ggccaggatga aacttgctga ctttggctct 600
gcttccatgg catcacctgc caattccttt gtgggaacgc cgtattggat ggccccagaa 660
gtaattttag ccatggatga aggacaatat gatggcaaag tagatgtgtg gtctcttgga 720
50 ataacatgta ttgaactagc ggaaaggaag cctcctttat ttaatatgaa tgcaatgagt 780
gccttatatc acatagccca aatgaatcc cctacactac agtctaata gaatggtctgat 840
55 tattttcgca actttgtaga ttcttgctc cagaaaatcc ctcaagatcg acctacatca 900
gaggaacttt taaagcacat attgttctt cgggagcgcc ctgaaaccgt gttaatatagat 960

```

ctcattcaga ggacaaagga tgcagtaaga gagctggaca atctgcagta tcgaaagatg 1020
5 aagaaactcc ttttccagga ggcacataat ggaccagcag tagaagcaca ggaagaagaa 1080
gaggaacaag atcatggtgt tggccggaca ggaacagtta atagtgttgg aagtaatcaa 1140
tccattccca gcatgtccat cagtgccagc agccaaagca gtagtgtaa cagtcttcca 1200
10 gatgtctcag atgacaagag tgagctagac atgatggagg gagaccacac agtgatgtct 1260
aacagttctg ttatccattt aaaaccagag gaagaaaatt acagagaaga gggagatcct 1320
agaacaagag catcagatcc acaatctcca cccaagtat ctctgcacaa atcacactat 1380
15 cgtaatcgag aacactttgc tactatacgg acagcatcac tggttacgag gcaaatgcaa 1440
gaacatgagc aggactctga gcttagagaa caaatgtctg gctataagcg aatgaggcga 1500
20 caacatcaaa agcaactgat gactctggaa aacaagctaa aggctgagat ggatgaacat 1560
cgcttcagat tagacaaaga tcttgaaact cagcgtaaca attttgctgc agaaatggag 1620
aaacttatca agaaacacca ggctgctatg gagaaagagg cttaaagtat gtccaatgaa 1680
25 gagaaaaaat ttcagcaaca tattcaggcc caacagaaga aagaactgaa tagttttctc 1740
gagtcccaga aaagagagta taaacttcga aaagagcagc ttaaagagga gctaaatgaa 1800
30 aaccagagta cccccaaaa agaaaaacag gagtggcttt caaagcagaa ggagaatata 1860
cagcatttcc aagcagaaga agaagctaac ctctctcgac gtcaaagaca atacctagag 1920
ctggaatgcc gtcgcttcaa gagaagaatg ttacttgggc gtcataactt agagcaggac 1980
35 cttgtcaggg aggagttaa caaaagacag actcagaagg acttagagca tgccatgcta 2040
ctccgacagc atgaatctat gcaagaactg gagttccgcc acctcaacac aattcagaag 2100
40 atgcgctgtg agttgatcag attacagcat caaactgagc tcactaacca gctggaatat 2160
aataagcgaa gagaacgaga actaagacga aagcatgtca tggaggttcg acaacagcct 2220
aagagtttga agtctaaaga actccaaata aaaaagcagt ttcaggatac ctgcaaaatc 2280
45 caaaccagac agtacaaagc attaagaaat cacctgctgg agactacacc aaagagttag 2340
cacaaagctg ttctgaaacg gctcaaggag gaacagaccc ggaaattagc tatcttggt 2400
50 gagcagtatg atcacagcat taatgaaatg ctctccacac aagccctgcg tttggatgaa 2460
gcacaggaag cagagtgccg ggttttgaag atgcagctgc agcaggaact ggagctgttg 2520
aatgcgtatc agagcaaaat caagatgcaa gctgaggcac aacatgatcg agagcttcgc 2580
55 gagcttgaac agaggggtctc cctccggagg gcactcttag aacaaaagat tgaagaagag 2640

```

    atgttggtt tgcagaatga gcgcacagaa cgaatacgaa gcctgttgga acgtcaagcc 2700
    agagagattg aagcttttga ctctgaaagc atgagactag gtttttagtaa tatggtgctt 2760
5   tctaattctct cccctgaggc attcagccac agctacccgg gagcttctgg ttggtcacac 2820
    aaccttactg ggggtccagg acctcactgg ggtcatccca tgggtggccc accacaagct 2880
    tggggccatc caatgcaagg tggaccccag ccatggggtc acccttcagg gccaatgcaa 2940
10  ggggtacctc gaggtagcag tatgggagtc cgcaatagcc cccaagctct gaggcggaca 3000
    gcttctgggg gacggacgga gcagggcatg agcagaagca cgagtgtcac ttcacaaata 3060
    tccaatgggt cacacatgtc ttatacataa cttaataatt gagagtggca attccgctgg 3120
    agctgtctgc caaaagaaac tgccacaga catcatcaca gcagcctcct cacttgggta 3180
    ctacagtgtg gaagctgagt gcatatggta tattttattc atttttgtaa agcgttctgt 3240
20  tttngtttta ctaattggga tgncatagta cttggctncc cgggt 3285

    <210> 10
25  <211> 1005
    <212> PRT
30  <213> Homo sapiens

    <300>
35  <308> SpTrEMBL
    <309> 2000-10-01
    <313> (1)..(1005)
40

    <400> 10
    Leu Leu Ser Arg Met Pro Ser Thr Asn Arg Ala Gly Ser Leu Lys Asp
45  1          5          10          15
    Pro Glu Ile Ala Glu Leu Phe Phe Lys Glu Asp Pro Glu Lys Leu Phe
          20          25          30
50  Thr Asp Leu Arg Glu Ile Gly His Gly Ser Phe Gly Ala Val Tyr Phe
          35          40          45
    Ala Arg Asp Val Arg Thr Asn Glu Val Val Ala Ile Lys Lys Met Ser
          50          55          60
55  Tyr Ser Gly Lys Gln Ser Thr Glu Lys Trp Gln Asp Ile Ile Lys Glu
    65          70          75          80

```

	Val	Lys	Phe	Leu	Gln	Arg	Ile	Lys	His	Pro	Asn	Ser	Ile	Glu	Tyr	Lys	
					85					90					95		
5	Gly	Cys	Tyr	Leu	Arg	Glu	His	Thr	Ala	Trp	Leu	Val	Met	Glu	Tyr	Cys	
				100					105					110			
	Leu	Gly	Ser	Ala	Ser	Asp	Leu	Leu	Glu	Val	His	Lys	Lys	Pro	Leu	Gln	
			115					120					125				
10	Glu	Val	Glu	Ile	Ala	Ala	Ile	Thr	His	Gly	Ala	Leu	Gln	Gly	Leu	Ala	
		130					135					140					
	Tyr	Leu	His	Ser	His	Thr	Met	Ile	His	Arg	Asp	Ile	Lys	Ala	Gly	Asn	
15	145					150					155					160	
	Ile	Leu	Leu	Thr	Glu	Pro	Gly	Gln	Val	Lys	Leu	Ala	Asp	Phe	Gly	Ser	
				165						170					175		
20	Ala	Ser	Met	Ala	Ser	Pro	Ala	Asn	Ser	Phe	Val	Gly	Thr	Pro	Tyr	Trp	
				180					185					190			
	Met	Ala	Pro	Glu	Val	Ile	Leu	Ala	Met	Asp	Glu	Gly	Gln	Tyr	Asp	Gly	
			195					200					205				
25	Lys	Val	Asp	Val	Trp	Ser	Leu	Gly	Ile	Thr	Cys	Ile	Glu	Leu	Ala	Glu	
		210					215					220					
	Arg	Lys	Pro	Pro	Leu	Phe	Asn	Met	Asn	Ala	Met	Ser	Ala	Leu	Tyr	His	
30	225					230					235					240	
	Ile	Ala	Gln	Asn	Glu	Ser	Pro	Thr	Leu	Gln	Ser	Asn	Glu	Trp	Ser	Asp	
				245						250					255		
35	Tyr	Phe	Arg	Asn	Phe	Val	Asp	Ser	Cys	Leu	Gln	Lys	Ile	Pro	Gln	Asp	
			260						265					270			
	Arg	Pro	Thr	Ser	Glu	Glu	Leu	Leu	Lys	His	Ile	Phe	Val	Leu	Arg	Glu	
			275					280					285				
40	Arg	Pro	Glu	Thr	Val	Leu	Ile	Asp	Leu	Ile	Gln	Arg	Thr	Lys	Asp	Ala	
		290					295					300					
	Val	Arg	Glu	Leu	Asp	Asn	Leu	Gln	Tyr	Arg	Lys	Met	Lys	Lys	Leu	Leu	
45	305					310					315					320	
	Phe	Gln	Glu	Ala	His	Asn	Gly	Pro	Ala	Val	Glu	Ala	Gln	Glu	Glu	Glu	
				325						330				335			
50	Glu	Glu	Gln	Asp	His	Gly	Val	Gly	Arg	Thr	Gly	Thr	Val	Asn	Ser	Val	
			340						345					350			
	Gly	Ser	Asn	Gln	Ser	Ile	Pro	Ser	Met	Ser	Ile	Ser	Ala	Ser	Ser	Gln	
			355					360					365				
55	Ser	Ser	Ser	Val	Asn	Ser	Leu	Pro	Asp	Val	Ser	Asp	Asp	Lys	Ser	Glu	
		370					375					380					

	Leu	Asp	Met	Met	Glu	Gly	Asp	His	Thr	Val	Met	Ser	Asn	Ser	Ser	Val	385	390	395	400
5	Ile	His	Leu	Lys	Pro	Glu	Glu	Glu	Asn	Tyr	Arg	Glu	Glu	Gly	Asp	Pro	405	410	415	
	Arg	Thr	Arg	Ala	Ser	Asp	Pro	Gln	Ser	Pro	Pro	Gln	Val	Ser	Arg	His	420	425	430	
10	Lys	Ser	His	Tyr	Arg	Asn	Arg	Glu	His	Phe	Ala	Thr	Ile	Arg	Thr	Ala	435	440	445	
	Ser	Leu	Val	Thr	Arg	Gln	Met	Gln	Glu	His	Glu	Gln	Asp	Ser	Glu	Leu	450	455	460	
15	Arg	Glu	Gln	Met	Ser	Gly	Tyr	Lys	Arg	Met	Arg	Arg	Gln	His	Gln	Lys	465	470	475	480
20	Gln	Leu	Met	Thr	Leu	Glu	Asn	Lys	Leu	Lys	Ala	Glu	Met	Asp	Glu	His	485	490	495	
	Arg	Leu	Arg	Leu	Asp	Lys	Asp	Leu	Glu	Thr	Gln	Arg	Asn	Asn	Phe	Ala	500	505	510	
25	Ala	Glu	Met	Glu	Lys	Leu	Ile	Lys	Lys	His	Gln	Ala	Ala	Met	Glu	Lys	515	520	525	
30	Glu	Ala	Lys	Val	Met	Ser	Asn	Glu	Glu	Lys	Lys	Phe	Gln	Gln	His	Ile	530	535	540	
	Gln	Ala	Gln	Gln	Lys	Lys	Glu	Leu	Asn	Ser	Phe	Leu	Glu	Ser	Gln	Lys	545	550	555	560
35	Arg	Glu	Tyr	Lys	Leu	Arg	Lys	Glu	Gln	Leu	Lys	Glu	Glu	Leu	Asn	Glu	565	570	575	
	Asn	Gln	Ser	Thr	Pro	Lys	Lys	Glu	Lys	Gln	Glu	Trp	Leu	Ser	Lys	Gln	580	585	590	
40	Lys	Glu	Asn	Ile	Gln	His	Phe	Gln	Ala	Glu	Glu	Glu	Ala	Asn	Leu	Leu	595	600	605	
45	Arg	Arg	Gln	Arg	Gln	Tyr	Leu	Glu	Leu	Glu	Cys	Arg	Arg	Phe	Lys	Arg	610	615	620	
	Arg	Met	Leu	Leu	Gly	Arg	His	Asn	Leu	Glu	Gln	Asp	Leu	Val	Arg	Glu	625	630	635	640
50	Glu	Leu	Asn	Lys	Arg	Gln	Thr	Gln	Lys	Asp	Leu	Glu	His	Ala	Met	Leu	645	650	655	
	Leu	Arg	Gln	His	Glu	Ser	Met	Gln	Glu	Leu	Glu	Phe	Arg	His	Leu	Asn	660	665	670	
55	Thr	Ile	Gln	Lys	Met	Arg	Cys	Glu	Leu	Ile	Arg	Leu	Gln	His	Gln	Thr	675	680	685	

Glu Leu Thr Asn Gln Leu Glu Tyr Asn Lys Arg Arg Glu Arg Glu Leu
 690 695 700
 5 Arg Arg Lys His Val Met Glu Val Arg Gln Gln Pro Lys Ser Leu Lys
 705 710 715 720
 Ser Lys Glu Leu Gln Ile Lys Lys Gln Phe Gln Asp Thr Cys Lys Ile
 725 730 735
 10 Gln Thr Arg Gln Tyr Lys Ala Leu Arg Asn His Leu Leu Glu Thr Thr
 740 745 750
 Pro Lys Ser Glu His Lys Ala Val Leu Lys Arg Leu Lys Glu Glu Gln
 15 755 760 765
 Thr Arg Lys Leu Ala Ile Leu Ala Glu Gln Tyr Asp His Ser Ile Asn
 770 775 780
 20 Glu Met Leu Ser Thr Gln Ala Leu Arg Leu Asp Glu Ala Gln Glu Ala
 785 790 795 800
 Glu Cys Gln Val Leu Lys Met Gln Leu Gln Gln Glu Leu Glu Leu Leu
 805 810 815
 25 Asn Ala Tyr Gln Ser Lys Ile Lys Met Gln Ala Glu Ala Gln His Asp
 820 825 830
 Arg Glu Leu Arg Glu Leu Glu Gln Arg Val Ser Leu Arg Arg Ala Leu
 30 835 840 845
 Leu Glu Gln Lys Ile Glu Glu Glu Met Leu Ala Leu Gln Asn Glu Arg
 850 855 860
 35 Thr Glu Arg Ile Arg Ser Leu Leu Glu Arg Gln Ala Arg Glu Ile Glu
 865 870 875 880
 Ala Phe Asp Ser Glu Ser Met Arg Leu Gly Phe Ser Asn Met Val Leu
 885 890 895
 40 Ser Asn Leu Ser Pro Glu Ala Phe Ser His Ser Tyr Pro Gly Ala Ser
 900 905 910
 Gly Trp Ser His Asn Pro Thr Gly Gly Pro Gly Pro His Trp Gly His
 45 915 920 925
 Pro Met Gly Gly Pro Pro Gln Ala Trp Gly His Pro Met Gln Gly Gly
 930 935 940
 50 Pro Gln Pro Trp Gly His Pro Ser Gly Pro Met Gln Gly Val Pro Arg
 945 950 955 960
 Gly Ser Ser Met Gly Val Arg Asn Ser Pro Gln Ala Leu Arg Arg Thr
 965 970 975
 55 Ala Ser Gly Gly Arg Thr Glu Gln Gly Met Ser Arg Ser Thr Ser Val
 980 985 990

Thr Ser Gln Ile Ser Asn Gly Ser His Met Ser Tyr Thr
 995 1000 1005

5 <210> 11
 <211> 1576
 10 <212> DNA
 <213> artificial sequence
 15 <220>
 <223> RNAi fragment T17E9.1a (kin-18)
 <400> 11
 20 cgaaaaccag cagaagagcg aatgtcagct gaagaatgct ttcgacatcc attcattcaa 60
 cggctctcgcc catcagacac aattcaggaa ctcatcaga gaacgaaaaa tatgggtatta 120
 gagttggata attttcaata caaaaagatg agaaaactca tgtatttgga tgaaacagaa 180
 25 ggaaaagaag gaagtgaagg aaatggagca tctgatgatt tagattttca tggaaatgaa 240
 gctaattcaa ttggaagagg tagtttttaa aattcaaagt gaaaatatta atatcttgga 300
 30 ataattttta taatattgct ttaaaccctc agcttttttt tgcagactct atcccttagt 360
 tgttcgtttt ccatctattc tcgttttcag caggagattc tgcgtcatct cgaagtgctt 420
 ctcttacttc tttccgatca atgcagagta gtggaggagc tggctcttta gtgtccacca 480
 35 atacgacggg tgctatggat aatgtgcatg gtactgtact gttttttttg ttttaggaat 540
 ggctttatta tttcctgcaa agttcaaaaa ttccatttat tttagttttt ctctcgaaat 600
 40 tcatcgcgca acattgagaa tctttcaaaa ttttcaggat cctctggata cggtaatgga 660
 agtagttcga cgacgagctc cgcacgccgc cgtcctcaa ttccttcgca aatgctctct 720
 tctacatcaa cgtctggtgt tggaaactatg ccgagtcatt gatcagttgg agcatcgatt 780
 45 acggcgatcg cagtcaatcc aacaccgtct ccttcagaac ctatcccaac atcacaacca 840
 acatcgaaat cagaatcatc ttctatactc gaaactgcac acgatgatcc tttggacacg 900
 50 tcgatacgtg ctccagtga agacttgcat atgccgcac gagcagtcaa ggaacgaata 960
 gccacgttgc aaaatcacia attcgcgacg cttcgttccc agagaataat caatcaggaa 1020
 caagaagaat atacgaaaga gaacaatatg tatgagcaaa tgagcaagta caagcatcta 1080
 55 cgacaagcac atcaciaaga gctccaacia tttgaagaac gatgtgcatt agatagagag 1140

caactgcgtg tgaaaatgga tcgagaactc gaacaattga caacgacata ctcgaaagaa 1200
aagatgagag tgaggtgttc acagaataat gaactagaca aacggaaaaa agatatcgaa 1260
5 gatggggaga aaaagatgaa aaagacgaaa aatagtcaaa atcagcagca gatgaaactg 1320
tattcagcgc aacaattgaa agaatacaag tataacaagg aggcacagaa aacagtgaga 1380
attcactttt atttgatttc tgtaaagaaa ttatacgaaa tttagacttt ataaattttt 1440
10 aaatatgaaa gttctggtca ctttttcagc tgcttctcca cttttttcaa agtttattat 1500
ttagtcttga ataatttttt aaaaaatgtc ctaaaccaag aattttcagc gattacgaag 1560
15 tctgaacatg cctcga 1576

<210> 12

20 <211> 20

<212> DNA

25 <213> artificial sequence

<220>

30 <223> primer

<400> 12

cgaaaaccag cagaagagcg 20

35 <210> 13

<211> 21

40 <212> DNA

<213> artificial sequence

<220>

45 <223> primer

<400> 13

50 tcgaggcatg ttcagacttc g 21